

New England Biolabs Certificate of Analysis

Product Name: T4 DNA Ligase
Catalog Number: M0202L
Concentration: 400,000 U/ml
Unit Definition: One unit is defined as the amount of enzyme required to give 50% ligation of 6 µg of Lambda-HindIII DNA in 30 minutes at 16°C in a total reaction volume of 20 µl.
Packaging Lot Number: 10181279
Expiration Date: 09/2024
Storage Temperature: -20°C
Storage Conditions: 10 mM Tris-HCl , 50 mM KCl , 1 mM DTT , 0.1 mM EDTA , 50 % Glycerol, (pH 7.4 @ 25°C)
Specification Version: PS-M0202S/L v1.0

T4 DNA Ligase Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
M0202LVIAL	T4 DNA Ligase	10161326	Pass
B0202SVIAL	T4 DNA Ligase Reaction Buffer	10153865	Pass


Assay Name/Specification	Lot # 10181279
<p>DNase Activity (Labeled Oligo, 3' extension) A 50 µl reaction in CutSmart® Buffer containing a 20 nM solution of a fluorescent labeled double-stranded oligonucleotide containing a 3' extension and a minimum of 10,000 units of T4 DNA Ligase incubated for 16 hours at 37°C yields <5% degradation as determined by capillary electrophoresis.</p>	Pass
<p>DNase Activity (Labeled Oligo, 5' extension) A 50 µl reaction in CutSmart® Buffer containing a 20 nM solution of a fluorescent labeled double-stranded oligonucleotide containing a 5' extension and a minimum of 10,000 units of T4 DNA Ligase incubated for 16 hours at 37°C yields <5% degradation as determined by capillary electrophoresis.</p>	Pass
<p>Double Stranded DNase Activity (Labeled Oligo) A 50 µl reaction in CutSmart® Buffer containing a 20 nM solution of a fluorescent labeled double-stranded oligonucleotide containing a blunt end and a minimum of 10,000 units of T4 DNA Ligase incubated for 16 hours at 37°C yields <5% degradation as determined by capillary electrophoresis.</p>	Pass

Assay Name/Specification	Lot # 10181279
<p>Endonuclease Activity (Nicking) A 50 µl reaction in NEBuffer 1 containing 1 µg of supercoiled PhiX174 DNA and a minimum of 2000 units of T4 DNA Ligase incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.</p>	Pass
<p>Exonuclease Activity (Radioactivity Release) A 50 µl reaction in NEBuffer 1 containing 1 µg of a mixture of single and double-stranded [³H] E. coli DNA and a minimum of 2000 units of T4 DNA Ligase incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.</p>	Pass
<p>Ligation and Recutting (Terminal Integrity, Digested DNA) A 20 µl reaction in 1X T4 DNA Ligase Reaction Buffer containing 2 µg of Lambda DNA-HindIII Digest and a minimum of 4000 units of T4 DNA Ligase incubated for 16 hours at 37°C results in >95% ligation of the DNA fragments as determined by agarose gel electrophoresis. Of these ligated fragments, >95% can be recut with HindIII.</p>	Pass
<p>Non-Specific DNase Activity (16 Hour) A 50 µl reaction in NEBuffer 1 containing 1 µg of CIP-treated Lambda-HindIII DNA and a minimum of 2000 units of T4 DNA Ligase incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.</p>	Pass
<p>Protein Concentration (A280) The concentration of T4 DNA Ligase is 0.4 mg/ml +/- 10% as determined by UV absorption at 280 nm. Protein concentration is determined by the Pace method using the extinction coefficient of 57,675 and molecular weight of 55,292 daltons for T4 DNA Ligase (Pace, C.N. et al. (1995) Protein Sci., 4, 2411-2423).</p>	Pass
<p>Protein Purity Assay (SDS-PAGE) T4 DNA Ligase is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.</p>	Pass
<p>RNase Activity (Extended Digestion) A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 µl of T4 DNA Ligase is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.</p>	Pass
<p>Single Stranded DNase Activity (FAM-Labeled Oligo) A 50 µl reaction in CutSmart® Buffer containing a 20 nM solution of a fluorescent internal labeled oligonucleotide and a minimum of 10,000 units of T4 DNA Ligase incubated for 16 hours at 37°C yields <5% degradation as determined by capillary electrophoresis.</p>	Pass

Assay Name/Specification	Lot # 10181279
<p>qPCR DNA Contamination (E. coli Genomic) A minimum of 2000 units of T4 DNA Ligase is screened for the presence of E. coli genomic DNA using SYBR® Green qPCR with primers specific for the E. coli 16S rRNA locus. Results are quantified using a standard curve generated from purified E. coli genomic DNA. The measured level of E. coli genomic DNA contamination is ≤ 1 E. coli genome.</p>	<p>Pass</p>

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.



Mary Lorenzen
Production Scientist
02 Sep 2022



Josh Hersey
Packaging Quality Control Inspector
01 Mar 2023