

www.neb.com info@neb.com



New England Biolabs Certificate of Analysis

Product Name: AMV Reverse Transcriptase

Catalog Number: M0277L
Concentration: 10,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme required to incorporate

1 nmol of dTTP into an acid-insoluble form in 10 minutes at 37°C.

Packaging Lot Number: 10226332
Expiration Date: 02/2026
Storage Temperature: -20°C

Storage Conditions: 200 mM KPO4, 2 mM DTT, 0.2 % Triton®X-100, 50 % Glycerol, (pH 7.2 @

25°C)

Specification Version: PS-M0277S/L v2.0

AMV Reverse Transcriptase Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
M0277LVIAL	AMV Reverse Transcriptase	10226330	Pass	
B0277AVIAL	AMV Reverse Transcriptase Reaction Buffer	10229630	Pass	

Assay Name/Specification	Lot # 10226332
Endonuclease Activity (Nicking) A 50 μl reaction in AMV Reverse Transcriptase Reaction Buffer containing 1 μg of supercoiled PhiX174 DNA and a minimum of 50 units of AMV Reverse Transcriptase incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
Exonuclease Activity (Radioactivity Release) A 50 µl reaction in AMV Reverse Transcriptase Reaction Buffer containing 1 µg of a mixture of single and double-stranded [³H] E. coli DNA and a minimum of 50 units of AMV Reverse Transcriptase incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass
Non-Specific DNase Activity (16 Hour) A 50 µl reaction in NEBuffer 2 containing 1 µg of T3 or T7 DNA in addition to a reaction containing Lambda-HindIII DNA and a minimum of 10 units of AMV Reverse Transcriptase incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass
RNAse Activity Assay (4 Hour Digestion)	Pass



M0277L / Lot: 10226332

Page 1 of 2



Assay Name/Specification	Lot # 10226332
A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA	
and a minimum of 1 µl of AMV Reverse Transcriptase is incubated at 37°C. After	
incubation for 4 hours, >90% of the substrate RNA remains intact as determined by	
gel electrophoresis using fluorescent detection.	

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Lea Antonopoulos

Michael Tonello

Packaging Quality Control Inspector

11 Apr 2024