

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

Product Name:	RNase Inhibitor, Murine
Catalog #:	M0314S/L
Concentration:	40,000 units/ml
Unit Definition:	One unit is defined as the amount of Murine RNase Inhibitor required to inhibit the activity of 5ng of RNase A by 50%. Activity is measured by the inhibition of hydrolysis of cytidine 2', 3'-cyclic monophosphate by RNase A.
Lot #:	0071405
Assay Date:	05/2014
Expiration Date:	5/2016
Storage Temp:	-20 °C
Storage Conditions:	50 mM KCl , 20 mM HEPES (pH 7.6), 8 mM DTT , 50 % Glycerol
Specification Version:	<i>PS-M0314S/L v1.0</i>
Effective Date:	15 May 2014

Assay Name/Specification (minimum release criteria)	Lot #0071405
<b>Endonuclease Activity (Nicking)</b> - A 50 µl reaction in NEBuffer 4 containing 1 µg of supercoiled PhiX174 DNA and a minimum of 40 units of RNase Inhibitor, Murine incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
<b>Exonuclease Activity (Radioactivity Release)</b> - A 50 $\mu$ l reaction in NEBuffer 4 containing 1 $\mu$ g of a mixture of single and double-stranded [ <sup>3</sup> H] <i>E. coli</i> DNA and a minimum of 200 units of RNase Inhibitor, Murine incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass
<b>Latent RNase Activity (Extended Digest)</b> - A 10 $\mu$ l reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 40 units of heat inactivated RNase Inhibitor, Murine is incubated at 37°C. After incubation for 4 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.	Pass
<b>Protein Purity Assay (SDS-PAGE)</b> - RNase Inhibitor, Murine is $\geq$ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	Pass
<b>RNase Activity (Extended Digestion)</b> - A 10 $\mu$ l reaction in NEBuffer 4 containing 40 ng of a 300 base single- stranded RNA and a minimum of 40 units of RNase Inhibitor, Murine is incubated at 37°C. After incubation for 4 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.	Pass

otunon

Authorized by Derek Robinson 15 May 2014



Honexian Gue

Inspected by Dongxian Yue 15 May 2014