

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name: SplintR® Ligase

Catalog Number: M0375L
Concentration: 25,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme needed to ligate (to 50%

completion) 2 picomoles of a tripartite FAM-labeled DNA:RNA hybrid substrate in a 20 µl reaction at 25°C in 15 minutes in 1X SplintR®

Ligase Reaction Buffer.

Packaging Lot Number: 10133511
Expiration Date: 12/2023
Storage Temperature: -20°C

Storage Conditions: 10 mM Tris-HCI, 300 mM NaCI, 1 mM DTT, 0.1 mM EDTA, 50 %

Glycerol, (pH 7.4 @ 25°C)

Specification Version: PS-M0375S/L v1.0

SplintR® Ligase Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
M0375LVIAL	SplintR® Ligase	10133510	Pass	
B0375SVIAL	10X SplintR® Ligase Reaction Buffer	10106685	Pass	

Assay Name/Specification	Lot # 10133511
Protein Purity Assay (SDS-PAGE)	Pass
SplintR® Ligase is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	
RNase Activity (Extended Digestion)	Pass
A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA	
and a minimum of 25 units of SplintR® Ligase is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel	
electrophoresis using fluorescent detection.	
Exonuclease Activity (Radioactivity Release)	Pass
A 50 µl reaction in SplintR® Ligase Reaction Buffer containing 1 µg of a mixture of	
single and double-stranded [³ H] E. coli DNA and a minimum of 125 units of SplintR® Ligase incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	
Ligase incubated for 4 flours at 37 G feleases <0.1% of the total fadioactivity.	
Endonuclease Activity (Nicking)	Pass
A 50 μl reaction in SplintR® Ligase Reaction Buffer containing 1 μg of supercoiled	



M0375L / Lot: 10133511

Page 1 of 2



Assay Name/Specification	Lot # 10133511
PhiX174 DNA and a minimum of 125 units of SplintR® Ligase incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Bhairavi Jani Production Scientist 07 Feb 2022 Michael Tonello Packaging Quality Control Inspector 07 Feb 2022

