

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

Product Name: SplintR® Ligase

Catalog Number: M0375S
Concentration: 25,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme needed to ligate (to 50%

completion) 2 picomoles of a tripartite FAM-labeled DNA:RNA hybrid substrate in a 20 µl reaction at 25°C in 15 minutes in 1X SplintR®

Ligase Reaction Buffer.

Packaging Lot Number: 10049232
Expiration Date: 07/2021
Storage Temperature: -20°C

Storage Conditions: 10 mM Tris-HCl , 300 mM NaCl , 1 mM DTT , 0.1 mM EDTA , 50 %

Glycerol, (pH 7.4 @ 25°C)

Specification Version: PS-M0375S/L v1.0

SplintR® Ligase Component List				
<b>NEB Part Number</b>	Component Description	Lot Number	Individual QC Result	
M0375SVIAL	SplintR® Ligase	10049233	Pass	
B0375SVIAL	10X SplintR® Ligase Reaction Buffer	10049234	Pass	

Assay Name/Specification	Lot # 10049232
Endonuclease Activity (Nicking)	Pass
A 50 μl reaction in SplintR® Ligase Reaction Buffer containing 1 μg of supercoiled	
PhiX174 DNA and a minimum of 125 units of SplintR® Ligase incubated for 4 hours at	
37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	
Exonuclease Activity (Radioactivity Release)	Pass
A 50 µl reaction in SplintR® Ligase Reaction Buffer containing 1 µg of a mixture of	
single and double-stranded [ ³H] E. coli DNA and a minimum of 125 units of SplintR® Ligase incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	
Ligase incubated for 4 flours at 57 C feleases <0.1% of the total fadioactivity.	
Protein Purity Assay (SDS-PAGE)	Pass
SplintR® Ligase is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie	
Blue detection.	
RNase Activity (Extended Digestion)	Pass
A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA	



M0375S / Lot: 10049232

Page 1 of 2

Assay Name/Specification	Lot # 10049232
and a minimum of 25 units of SplintR® Ligase is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.	

This product has been tested and shown to be in compliance with all specifications.

Do Mu

Production Scientist 15 Jul 2019 Michael Tonello

Packaging Quality Control Inspector

13 Dec 2019

