

New England Biolabs Certificate of Analysis

Product Name: dATP Solution

 Catalog #:
 N0440S

 Concentration:
 100 mM

 Unit Definition:
 N/A

 Lot #:
 0621801

 Assay Date:
 01/2018

 Expiration Date:
 1/2020

 Storage Temp:
 -20°C

Storage Conditions: Supplied in Ultrapure water as a sodium salt (pH 7.5)

Specification Version: PS-N0440S v1.0 Effective Date: 24 Jan 2018

Assay Name/Specification (minimum release criteria)	Lot #0621801
Endonuclease Activity (Nicking) - A 50 μ l reaction in NEBuffer 2 containing 1 μ g of supercoiled PhiX174 DNA and a minimum of 1 μ l of dATP Solution incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
Non-Specific DNase Activity (16 Hour) - A 50 µl reaction in NEBuffer 2 containing 1 µg of T3 DNA in addition to a reaction containing Lambda-HindIII DNA and a minimum of 10 µl of dATP Solution incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass
PCR Amplification (0.5 kb Lambda, dNTPs) - A 50 μl reaction in ThermoPol® Reaction Buffer in the presence of 200 μM dATP, dGTP, dCTP, and dTTP and 0.5 μM primers containing 1 ng Lambda DNA with 1.25 units of <i>Taq</i> DNA Polymerase for 25 cycles of PCR amplification results in the expected 0.5 kb product.	Pass
PCR Amplification (2.0 kb Lambda, dNTPs) - A 50 μl reaction in ThermoPol® Reaction Buffer in the presence of 200 μM dATP, dGTP, dCTP, and dTTP and 0.5 μM primers containing 1 ng Lambda DNA with 1.25 units of <i>Taq</i> DNA Polymerase for 25 cycles of PCR amplification results in the expected 2.0 kb product.	Pass
PCR Amplification (5.0 kb Lambda, dNTPs) - A 50 μl reaction in ThermoPol® Reaction Buffer in the presence of 200 μM dATP, dGTP, dCTP, and dTTP and 0.5 μM primers containing 1 ng Lambda DNA with 1.25 units of <i>Taq</i> DNA Polymerase for 25 cycles of PCR amplification results in the expected 5.0 kb product.	Pass
Phosphatase Activity (pNPP) - A 200 μl reaction in 1M Diethanolamine, pH 9.8, 0.5 mM MgCl ₂ containing 2.5 mM <i>p</i> -Nitrophenyl Phosphate (pNPP) and a minimum of 4 μl dATP Solution incubated for 4 hours at 37°C yields <0.0001 unit of alkaline phosphatase activity as determined by spectrophotometric analysis.	Pass









240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Assay Name/Specification (minimum release criteria)	Lot #0621801
Physical Purity (HPLC) - dATP Solution is ≥ 99% pure as determined by HPLC analysis.	Pass
RNase Activity (Extended Digestion) - A 10 μl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 μl of dATP Solution is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.	Pass

Authorized by Lynne Apone 24 Jan 2018







Inspected by
Tony Spear-Alfonso
02 Feb 2018