

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name: Bacteroides Heparinase II

Catalog Number: P0736L
Concentration: 4,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme that will liberate 1.0

µmol unsaturated oligosaccharides from porcine mucosal heparin per

minute at 30°C and pH 7.0 in a total reaction volume of 100 μl.

Packaging Lot Number: 10052582
Expiration Date: 09/2020
Storage Temperature: -80°C

Storage Conditions: 100 mM NaCl, 20 mM Tris-HCl, 1 mM EDTA, 5 mM CaCl2, (pH 7.5 @ 25°C)

Specification Version: PS-P0736S/L v1.0

Bacteroides Heparinase II Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
P0736LVIAL	Bacteroides Heparinase II	10052581	Pass	
B0735SVIAL	Bacteroides Heparinase Reaction Buffer (10X)	10052584	Pass	

Assay Name/Specification	Lot # 10052582
Protein Purity Assay (SDS-PAGE) Bacteroides Heparinase II is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	Pass
Protease Activity (SDS-PAGE) A 20 µl reaction in 1X Heparinase Reaction Buffer containing 24 µg of a standard mixture of proteins and a minimum of 20 units of Bacteroides Heparinase II incubated for 20 hours at 37°C, results in no detectable degradation of the protein mixture as determined by SDS-PAGE with Coomassie Blue detection.	Pass
Glycosidase Activity (β1-4 Galactosidase) A 10 μl reaction in Heparinase Reaction Buffer containing 1 nM of fluorescently-labeled β-Galactosidase substrate (Galβ1-4GlcNAcβ1-3Galβ1-4Glc -AMC) and 8 units of Bacteroides Heparinase II incubated for 20 hours at 30°C results in no detectable activity as determined by thin layer chromatography.	Pass
Glycosidase Activity (β1-3 Galactosidase) A 10 μl reaction in Heparinase Reaction Buffer containing 1 nM of fluorescently-labeled β-Galactosidase substrate (Galβ1-3GlcNAcβ1-4Galβ1-4Glc-AMC)	Pass



P0736L / Lot: 10052582 Page 1 of 2

Assay Name/Specification	Lot # 10052582
and 8 units of Bacteroides Heparinase II incubated for 20 hours at 30°C results in no detectable activity as determined by thin layer chromatography.	
Glycosidase Activity (β-N-Acetylgalactosaminidase) A 10 μl reaction in Heparinase Reaction Buffer containing 1 nM of fluorescently-labeled β-N-Acetylgalactosaminidase substrate (GalNAcβ1-4Galβ1-4Glc-AMC) and 8 units of Bacteroides Heparinase II incubated for 20 hours at 30°C results in no detectable activity as determined by thin layer chromatography.	Pass
Glycosidase Activity (β-N-Acetylglucosaminidase) A 10 μl reaction in Heparinase Reaction Buffer containing 1 nM of fluorescently-labeled β-N-Acetylglucosaminidase substrate (GlcNAcβ1-4GlcNAcβ1-4GlcNAc-AMC) and 8 units of Bacteroides Heparinase II incubated for 20 hours at 30°C results in no detectable activity as determined by thin layer chromatography.	Pass
Sulfatase Activity (2-O) A 10 μl reaction in Heparinase Reaction Buffer containing 1 nM of fluorescently-labeled 2-O-Sulfatase substrate (ΔUA2S-(1-4)-GlcNS6S-AMC) and 8 units of Bacteroides Heparinase II incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	Pass
Sulfatase and Uronidase Activity (N,6-O) A 10 μl reaction in Heparinase Reaction Buffer containing 1 nM of fluorescently-labeled N,6-O-Sulfatase substrate (ΔUA-(1-4)-GlcNS6S-AMC) and 8 units of Bacteroides Heparinase II incubated for 20 hours at 30°C results in no detectable activity as determined by thin layer chromatography.	Pass

This product has been tested and shown to be in compliance with all specifications.

Brad Landgraf Production Scientist

09 Sep 2019

Josh Hersey

Packaging Quality Control Inspector

05 Nov 2019



P0736L / Lot: 10052582

Page 2 of 2