

*be* INSPIRED *drive* DISCOVERY *stay* GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

| Product Name:          | Random Primer Mix  |
|------------------------|--|
| Catalog Number:        | S1330S   |
| Concentration:         | 60 µM  |
| Packaging Lot Number:  | 10116777   |
| Expiration Date:       | 02/2024  |
| Storage Temperature:   | -20°C  |
| Specification Version: | PS-S1330S v2.0   |
| Composition (1X):      | 1 mM dATP, 1 mM dCTP, 1 mM dGTP, 1 mM dTTP, 35 μM Hexamers, 25 μM<br>dT(23)VN supplied in ultrapure water. |

| Random Primer Mix Component List |                       |            |                      |  |
|----------------------------------|-----------------------|------------|----------------------|--|
| NEB Part Number                  | Component Description | Lot Number | Individual QC Result |  |
| S1330SVIAL                       | Random Primer Mix     | 10099159   | Pass                 |  |

| Assay Name/Specification   | Lot # 10116777 |
|--|----------------|
| <b>Endonuclease Activity (Nicking)</b><br>A 25 $\mu$ I reaction in NEBuffer 2 containing 1 $\mu$ g of supercoiled PhiX174 DNA and a<br>minimum of 5 $\mu$ I of Random Primer Mix incubated for 4 hours at 37°C results in <10%<br>conversion to the nicked form as determined by agarose gel electrophoresis.  | Pass           |
| <b>Non-Specific DNase Activity (16 Hour)</b><br>A 50 µl reaction in NEBuffer 2 containing 1 µg of T3 or T7 DNA in addition to a<br>reaction containing Lambda-HindIII DNA and a minimum of 5 µl of Random Primer Mix<br>incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease<br>degradation as determined by agarose gel electrophoresis. | Pass           |
| <b>Phosphatase Activity (pNPP)</b><br>A 200 µl reaction in 1M Diethanolamine, pH 9.8, 0.5 mM MgCl2 containing 2.5 mM<br>p-Nitrophenyl Phosphate (pNPP) and a minimum of 20 µl of Random Primer Mix incubated<br>for 4 hours at 37°C yields <0.0001 unit of alkaline phosphatase activity as<br>determined by spectrophotometric analysis.                                | Pass           |
| <b>RNase Activity (Extended Digestion)</b><br>A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA<br>and a minimum of 1 µl of Random Primer Mix is incubated at 37°C. After incubation<br>for 16 hours, >90% of the substrate RNA remains intact as determined by gel<br>electrophoresis using fluorescent detection.                      | Pass           |





be INSPIRED drive DISCOVERY stay GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

vastie Vayanez

Christie Vazquez Production Scientist 27 Jul 2021

1 lon M

Mary Neal Packaging Quality Control Inspector 27 Jul 2021

