

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name: NEBuffer™ 3

Catalog Number: B7003S

Concentration: 10 X Concentrate

Packaging Lot Number: 10157349
Expiration Date: 04/2025
Storage Temperature: -20°C

Specification Version: PS-B7003S v1.0

Composition (1X): 100 mM NaCl, 50 mM Tris-HCl, 10 mM MgCl2, 1 mM DTT, (pH 7.9 @ 25°C)

| NEBuffer™ 3 Component List | | | | |
|----------------------------|-----------------------|------------|----------------------|--|
| NEB Part Number | Component Description | Lot Number | Individual QC Result | |
| B7003SVIAL | NEBuffer™ 3 | 10143290 | Pass | |

| Assay Name/Specification | Lot # 10157349 |
|---|----------------|
| RNase Activity (Buffer) A 10 µl reaction in 1X NEBuffer 3 containing 40 ng of a 300 base single-stranded RNA is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by fluorescent detection. | Pass |
| pH (buffers/solutions) The pH of 10X NEBuffer 3 is between pH 7.8 and 8.0 at 25°C. | Pass |
| Conductivity (buffers/solutions) The conductivity of 10X NEBuffer 3 is between 85 and 115 mS at 25°C. | Pass |
| Endonuclease Activity (Nicking, Buffer) A 50 μl reaction in 1X NEBuffer 3 containing 1 μg of supercoiled PhiX174 DNA incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis. | Pass |
| Non-Specific DNase Activity (16 hour, Buffer) A 50 µl reaction in 1X NEBuffer 3 containing 1 µg of PhiX174-HaeIII DNA incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis. | Pass |
| Functional Testing (Restriction Digest, BSA, Buffer) A 50 µl reaction in 1X NEBuffer 3 plus 100 µg/ml Bovine Serum Albumin containing 1 | Pass |



B7003S / Lot: 10157349

Page 1 of 2

| Assay Name/Specification | Lot # 10157349 |
|--|----------------|
| μg of pBC4 DNA and 1 unit of NotI incubated for 1 hour at 37°C results in complete digestion of the substrate DNA as determined by agarose gel electrophoresis. | |
| Functional Testing (Restriction Digest, BSA, Buffer) A 50 µl reaction in 1X NEBuffer 3 plus 100 µg/ml Bovine Serum Albumin containing 1 µg of Lambda DNA and 1 unit of Asel incubated for 1 hour at 37°C results in complete | Pass |

This product has been tested and shown to be in compliance with all specifications.

digestion of the substrate DNA as determined by agarose gel electrophoresis.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Ben Penta

Production Scientist

Bu Rear

20 Jul 2022

Erin Varney

Packaging Quality Control Inspector

20 Jul 2022



B7003S / Lot: 10157349

Page 2 of 2