

be INSPIRED *drive* DISCOVERY *stay* GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

| Product Name: | T7 RNA Polymerase |
|------------------------|---|
| Catalog Number: | M0251L |
| Concentration: | 50,000 U/ml |
| Unit Definition: | One unit is defined as the amount of enzyme that will incorporate 1 nmol ATP into acid-insoluble material in a total reaction volume of 50 μl in 1 hour at 37°C in 1X RNA Polymerase Reaction Buffer. |
| Lot Number: | 10035565 |
| Expiration Date: | 11/2020 |
| Storage Temperature: | -20°C |
| Storage Conditions: | 100 mM NaCl , 50 mM Tris-HCl (pH 7.9), 1 mM EDTA , 20 mM BME , 0.1 % Triton X-100 , 50 % Glycerol |
| Specification Version: | PS-M0251S/L v3.0 |

| T7 RNA Polymerase Component List | | | | |
|----------------------------------|------------------------|------------|----------------------|--|
| NEB Part Number | Component Description | Lot Number | Individual QC Result | |
| M0251LVIAL | T7 RNA Polymerase | 10026490 | Pass | |
| B9012SVIAL | RNAPol Reaction Buffer | 0181804 | Pass | |

| Assay Name/Specification | Lot # 10035565 |
|---|----------------|
| Endonuclease Activity (Nicking) A 50 µl reaction in RNAPol Reaction Buffer containing 1 µg of supercoiled PhiX174 DNA and a minimum of 150 units of T7 RNA Polymerase incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis. | Pass |
| Exonuclease Activity (Radioactivity Release) A 50 µl reaction in RNAPol Reaction Buffer containing 1 µg of a mixture of single and double-stranded [³ H] E. coli DNA and a minimum of 150 units of T7 RNA Polymerase incubated for 4 hours at 37°C releases <0.1% of the total radioactivity. | Pass |
| Non-Specific DNase Activity (16 Hour) A 50 µl reaction in RNAPol Reaction Buffer containing 1 µg of Lambda DNA and a minimum of 250 units of T7 RNA Polymerase incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis. | Pass |
| Promoter Specificity | Pass |





be INSPIRED drive DISCOVERY stay GENUINE

240 County Road Ipswich, MA 01938-2723

Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

| Assay Name/Specification | Lot # 10035565 |
|--|----------------|
| A 50 μ I reaction in RNAPol Reaction Buffer in the presence of 2 mM NTPs containing 1 μ g of Lambda DNA as a template and a minimum of 200 units of T7 RNA Polymerase incubated for 1 hour at 37°C results in <1.5% of the amount of product incorporated as compared to a control reaction using T7 DNA as a template. | |
| Protein Purity Assay (SDS-PAGE) T7 RNA Polymerase is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection. | Pass |
| RNase Activity (Extended Digestion) A 10 µl reaction in RNAPol Reaction Buffer containing 40 ng of a 300 base single-stranded RNA and a minimum of 50 units of T7 RNA Polymerase is incubated at 37°C. After incubation for 4 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection. | Pass |

This product has been tested and shown to be in compliance with all specifications.

ongeran he

Dongxian Yue Production Scientist 21 Nov 2018

Mich 11.

Michael Tonello Packaging Quality Control Inspector 13 Mar 2019

