

be INSPIRED drive DISCOVERY stay GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

| Product Name: | NEBNext® dsDNA Fragmentase® |
|------------------------|----------------------------------------------------------------------------------------------------------------|
| Catalog Number: | M0348S |
| Unit Definition: | |
| Lot Number: | 10027388 |
| Expiration Date: | 04/2020 |
| Storage Temperature: | -20°C |
| Storage Conditions: | 10 mM Tris-HCl, 50 mM NaCl, 0.1 mM EDTA, 50 % Glycerol, 0.15 % Triton®X-100, 200 μg/ml BSA, (pH 7.5 @ 25°C) |
| Specification Version: | PS-M0348S/L v1.0 |

| NEBNext® dsDNA Fragmentase® Component List | | | |
|--------------------------------------------|---------------------------------------------------|------------|----------------------|
| NEB Part Number | Component Description | Lot Number | Individual QC Result |
| M0348AVIAL | NEBNext® dsDNA Fragmentase® | 10025256 | Pass |
| B0511AVIAL | MgCl2 Solution (200 mM) | 10025258 | Pass |
| B0349AVIAL | NEBNext® dsDNA Fragmentase® Reaction Buffer v2 | 10025257 | Pass |

| Assay Name/Specification | Lot # 10027388 |
|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------|
| Phosphatase Activity (pNPP) A 200 µl reaction in 1M Diethanolamine, pH 9.8, 0.5 mM MgCl2 containing 2.5 mM p-Nitrophenyl Phosphate (pNPP) and a minimum of 10 µl of NEBNext® dsDNA Fragmentase® incubated for 4 hours at 37°C yields <0.00001 unit of alkaline phosphatase activity as determined by spectrophotometric analysis. | Pass |
| Protease Activity (SDS-PAGE) A 20 µl reaction in 1X NEBNext® dsDNA Fragmentase Reaction Buffer containing 24 µg of a standard mixture of proteins and a minimum of 10 µl of NEBNext® dsDNA Fragmentase® incubated for 20 hours at 37°C, results in no detectable degradation of the protein mixture as determined by SDS-PAGE with Coomassie Blue detection. | Pass |
| Protein Purity Assay (SDS-PAGE) NEBNext® dsDNA Fragmentase® is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection. | Pass |

This product has been tested and shown to be in compliance with all specifications.





be INSPIRED *drive* DISCOVERY *stay* GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

ina din

Melissa Arn Production Scientist 23 Oct 2018

Michae 711.

Michael Tonello Packaging Quality Control Inspector 26 Oct 2018

