

New England Biolabs Certificate of Analysis

Product Name: OneTaq[®] Hot Start Quick-Load[®] 2X Master Mix with GC Buffer
Catalog Number: M0489L
Concentration: 2 X Concentrate
Packaging Lot Number: 10105615
Expiration Date: 01/2023
Storage Temperature: -20°C
Specification Version: PS-M0489S/L v2.0
Composition (1X): 80 mM Tris-SO₄ (pH 9.2 @ 25°C), 20 mM (NH₄)₂SO₄, 2 mM MgSO₄, 0.2 mM dATP, 0.2 mM dCTP, 0.2 mM dGTP, 0.2 mM dTTP, 5 % Glycerol, 5 % DMSO, 0.06 % IGEPAL[®] CA-630, 0.05 % Tween[®] 20, 1 X Xylene cyanol, 1 X Tartrazine, 25 units/ml OneTaq[®] Hot Start DNA Polymerase

OneTaq [®] Hot Start Quick-Load [®] 2X Master Mix with GC Buffer Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
M0489SVIAL	OneTaq [®] Hot Start Quick-Load [®] 2X Master Mix with GC Buffer	10099130	Pass
B9026AVIAL	OneTaq [®] High GC Enhancer	10061963	Pass

Assay Name/Specification	Lot # 10105615
<p>Non-Specific DNase Activity (16 hour, Buffer) A 50 µl reaction in 1X OneTaq[®] Hot Start Quick-Load[®] Master Mix with GC Buffer containing 1 µg of T3 or T7 DNA in addition to a reaction containing Lambda-HindIII DNA incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.</p>	Pass
<p>RNase Activity (Extended Digestion) A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 µl of OneTaq[®] Hot Start Quick-Load[®] 2X Master Mix with GC Buffer is incubated at 37°C. After incubation for 4 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.</p>	Pass
<p>PCR Amplification (Buffer Dependent, >65% GC-rich, Master Mix) A 25 µl reaction in 1X OneTaq[®] Hot Start Quick-Load[®] Master Mix with GC Buffer and 0.2 µM primers containing 10 ng Human Genomic DNA for 30 cycles of PCR amplification results in the buffer-dependent production of the 737 bp product.</p>	Pass
<p>PCR Amplification (Hot Start 2 kb Lambda DNA)</p>	Pass


Assay Name/Specification	Lot # 10105615
<p>A 25 µl reaction in OneTaq[®] Standard Reaction Buffer in the presence of 200 µM dNTPs and 0.2 µM primers containing 10 pg Lambda DNA and 50 ng Human Genomic DNA with 0.625 units of OneTaq[®] Hot Start DNA Polymerase for 30 cycles of PCR amplification results in an increase in yield of the 2 kb Lambda product and a decrease in non-specific genomic bands when compared to a non-hot start control reaction.</p>	
<p>PCR Amplification (Enhancer Dependent, >70% GC-rich, Master Mix) A 25 µl reaction in 1X OneTaq[®] Hot Start Quick-Load[®] Master Mix with GC Buffer and 20% OneTaq[®] High GC Enhancer in the presence of 0.2 µM primers containing 10 ng Human Genomic DNA for 30 cycles of PCR amplification results in the enhancer-dependent production of the 627 bp product.</p>	Pass
<p>Inhibition of Primer Extension (Hot Start, Radioactivity Incorporation) A 50 µl primer extension assay in ThermoPol[®] Reaction Buffer in the presence of 200 µM dNTPs including [³H]-dTTP, containing 15 nM primed single-stranded M13mp18 with 2.5 units of OneTaq[®] Hot Start DNA Polymerase incubated for 16 hours at 25°C yields >95% inhibition when compared to a non-hot start control reaction.</p>	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.



Christie Vazquez
Production Scientist
22 Apr 2021



Michael Tonello
Packaging Quality Control Inspector
22 Apr 2021