

## New England Biolabs Certificate of Analysis

**Product Name:** SfcI  
**Catalog Number:** R0561L  
**Concentration:** 10,000 U/ml  
**Unit Definition:** One unit is defined as the amount of enzyme required to digest 1 µg of Lambda DNA in 1 hour at 37°C in a total reaction volume of 50 µl.  
**Packaging Lot Number:** 10159565  
**Expiration Date:** 08/2024  
**Storage Temperature:** -20°C  
**Storage Conditions:** 300 mM NaCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 500 µg/ml BSA  
**Specification Version:** PS-R0561S/L v1.0

| SfcI Component List |                       |            |                      |
|---------------------|-----------------------|------------|----------------------|
| NEB Part Number     | Component Description | Lot Number | Individual QC Result |
| R0561LVIAL          | SfcI                  | 10159564   | Pass                 |
| B6004SVIAL          | rCutSmart™ Buffer     | 10154051   | Pass                 |

| Assay Name/Specification  | Lot # 10159565 |
|---|----------------|
| <p><b>Ligation and Recutting (Terminal Integrity)</b><br/>           After a 2-fold over-digestion of Lambda DNA with SfcI, &gt;95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, &gt;95% can be recut with SfcI.</p>   | Pass           |
| <p><b>Exonuclease Activity (Radioactivity Release)</b><br/>           A 50 µl reaction in CutSmart™ Buffer containing 1 µg of a mixture of single and double-stranded [<sup>3</sup>H] E. coli DNA and a minimum of 50 units of SfcI incubated for 4 hours at 37°C releases &lt;0.1% of the total radioactivity.</p>   | Pass           |
| <p><b>Non-Specific DNase Activity (16 hour)</b><br/>           A 50 µl reaction in CutSmart™ Buffer containing 1 µg of Lambda DNA and a minimum of 10 Units of SfcI incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis. NOTE: although no nuclease degradation is detected under these conditions, extended incubations and/or high concentrations of this enzyme may result in star activity. See the product FAQ for recommended reaction conditions for this enzyme.</p> | Pass           |

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit [www.neb.com/trademarks](http://www.neb.com/trademarks) for additional information.



Penghua Zhang  
Production Scientist  
09 Sep 2022



Michael Tonello  
Packaging Quality Control Inspector  
09 Sep 2022