

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name: PaqCl®
Catalog Number: R0745L
Concentration: 10,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg

of Lambda DNA incubated for 1 hour at 37°C in a total reaction

volume of 50 μl.

Packaging Lot Number: 10171799
Expiration Date: 06/2024
Storage Temperature: -20°C

Storage Conditions: 300 mM NaCl, 10 mM Tris-HCl, 1 mM DTT, 0.1 mM EDTA, 50 %

Glycerol , 500 μg/ml BSA, (pH 7.4 @ 25°C)

Specification Version: PS-R0745S/L v1.0

PaqCI® Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
S0532LVIAL	PaqCI® Activator	10174169	Pass	
R0745LVIAL	PaqCl®	10171795	Pass	
B6004SVIAL	rCutSmart™ Buffer	10173161	Pass	

Assay Name/Specification	Lot # 10171799
Protein Purity Assay (SDS-PAGE)	Pass
PaqCl® is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	
Exonuclease Activity (Radioactivity Release)	Pass
A 50 ul reaction in CutSmart® Buffer containing 1 µg of a mixture of single and	
double-stranded [³H] E. coli DNA and a minimum of 100 units of PaqCI® incubated	
for 4 hours at 37°C releases <0.1% of the total radioactivity.	
Non-Specific DNase Activity (16 Hour)	Pass
A 50 µl reaction in CutSmart® Buffer containing 1 µg of Lambda DNA and a minimum of	
30 units of PaqCI® incubated for 16 hours at 37°C results in a DNA pattern free of	
detectable nuclease degradation as determined by agarose gel electrophoresis.	
Endonuclease Activity (Nicking)	Pass
A 50 µl reaction in CutSmart® Buffer containing 1 µg of supercoiled PhiX174 DNA and	
a minimum of 30 units of PaqCI® incubated for 4 hours at 37°C results in <20%	



R0745L / Lot: 10171799 Page 1 of 2

Assay Name/Specification	Lot # 10171799
conversion to the nicked form as determined by agarose gel electrophoresis.	
Ligation and Recutting (Terminal Integrity) After a 10-fold over-digestion of Lambda DNA with PaqCl®, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with PaqCl®.	Pass
Functional Testing (15 minute Digest) A 50 µl reaction in CutSmart® Buffer containing 1 µg of Lambda DNA and 1 µl of PaqCl® incubated for 15 minutes at 37°C results in complete digestion as determined by agarose gel electrophoresis.	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

YunJie Suń

Production Scientist

13 Dec 2022

Mary Negl

Packaging Quality Control Inspector

14 Dec 2022



R0745L / Lot: 10171799

Page 2 of 2