

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name: PaqCl®
Catalog Number: R0745L
Concentration: 10,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg

of Lambda DNA incubated for 1 hour at 37°C in a total reaction

volume of 50 μl.

Packaging Lot Number: 10177792
Expiration Date: 06/2024
Storage Temperature: -20°C

Storage Conditions: 300 mM NaCl, 10 mM Tris-HCl, 1 mM DTT, 0.1 mM EDTA, 50 %

Glycerol , 500 μg/ml BSA, (pH 7.4 @ 25°C)

Specification Version: PS-R0745S/L v1.0

PaqCI® Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
S0532LVIAL	PaqCI® Activator	10174169	Pass	
R0745LVIAL	PaqCI®	10177789	Pass	
B6004SVIAL	rCutSmart™ Buffer	10173663	Pass	

Assay Name/Specification	Lot # 10177792
Functional Testing (15 minute Digest)	Pass
A 50 μl reaction in CutSmart® Buffer containing 1 μg of Lambda DNA and 1 μl of	
PaqCI® incubated for 15 minutes at 37°C results in complete digestion as determined	
by agarose gel electrophoresis.	
Endonuclease Activity (Nicking)	Pass
A 50 µl reaction in CutSmart® Buffer containing 1 µg of supercoiled PhiX174 DNA and	
a minimum of 30 units of PaqCl® incubated for 4 hours at 37°C results in <20%	
conversion to the nicked form as determined by agarose gel electrophoresis.	
Protein Purity Assay (SDS-PAGE)	Pass
PaqCl® is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue	
detection.	
Ligation and Recutting (Terminal Integrity)	Pass
After a 10-fold over-digestion of Lambda DNA with PaqCl® , >95% of the DNA fragments	1 400
can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments,	



R0745L / Lot: 10177792 Page 1 of 2

Assay Name/Specification	Lot # 10177792
>95% can be recut with PaqCI®.	
Non-Specific DNase Activity (16 Hour) A 50 µl reaction in CutSmart® Buffer containing 1 µg of Lambda DNA and a minimum of 30 units of PaqCl® incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass
Exonuclease Activity (Radioactivity Release) A 50 ul reaction in CutSmart® Buffer containing 1 µg of a mixture of single and double-stranded [³H] E. coli DNA and a minimum of 100 units of PaqCl® incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

YunJie Suń

Production Scientist

13 Dec 2022

Michael Tonello

Packaging Quality Control Inspector

06 Feb 2023



R0745L / Lot: 10177792

Page 2 of 2