

## New England Biolabs Product Specification

<i>Product Name:</i>	<i>XhoI</i>
<i>Catalog #:</i>	<i>R0146S/L/E</i>
<i>Concentration:</i>	<i>20,000 units/ml</i>
<i>Unit Definition:</i>	<i>One unit is defined as the amount of enzyme required to digest 1 µg of Lambda DNA (HindIII digest) fragments in rCutSmart Buffer in 1 hour at 37°C in a total reaction volume of 50 µl.</i>
<i>Shelf Life:</i>	<i>24 months</i>
<i>Storage Temp:</i>	<i>-20°C</i>
<i>Storage Conditions:</i>	<i>10 mM Tris-HCl, 50 mM KCl, 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 200 µg/ml rAlbumin (pH 7.4 @ 25°C)</i>
<i>Specification Version:</i>	<i>PS-R0146S/L/E v3.0</i>
<i>Effective Date:</i>	<i>07 Oct 2022</i>

### Assay Name/Specification (minimum release criteria)

**Blue-White Screening (Terminal Integrity)** - A sample of Litmus 28i vector linearized with a 10-fold excess of XhoI, religated and transformed into an *E. coli* strain expressing the LacZ beta fragment gene results in <1% white colonies.

**Ligation and Recutting (Terminal Integrity)** - After a 10-fold over-digestion of pXba DNA with XhoI, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with XhoI.

**Protein Purity Assay (SDS-PAGE)** - XhoI is >95% pure as determined by SDS PAGE analysis using Coomassie Blue detection.

**Endonuclease Activity (Nicking)** - A 50 µl reaction in rCutSmart™ Buffer containing 1 µg of supercoiled pBR322 DNA and a minimum of 100 units of XhoI incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.

**Exonuclease Activity (Radioactivity Release)** - A 50 µl reaction in rCutSmart™ Buffer containing 1 µg of a mixture of single and double-stranded [<sup>3</sup>H] *E. coli* DNA and a minimum of 100 units of XhoI incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.

**Functional Testing (15 minute Digest)** - A 50 µl reaction in rCutSmart™ Buffer containing 1 µg of Lambda-HindIII DNA and 1 µl of XhoI incubated for 15 minutes at 37°C results in complete digestion as determined by agarose gel electrophoresis.

**Non-Specific DNase Activity (16 Hour)** - A 50 µl reaction in rCutSmart™ Buffer containing 1 µg of Lambda-HindIII DNA and a minimum of 100 units of XhoI incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.



---

## New England Biolabs Product Specification

<b>Assay Name/Specification (minimum release criteria)</b>
--

<b>qPCR DNA Contamination (<i>E. coli</i> Genomic)</b> - A minimum of 20 units of XhoI is screened for the presence of <i>E. coli</i> genomic DNA using SYBR <sup>®</sup> Green qPCR with primers specific for the <i>E. coli</i> 16S rRNA locus. Results are quantified using a standard curve generated from purified <i>E. coli</i> genomic DNA. The measured level of <i>E. coli</i> genomic DNA contamination is $\leq 1$ <i>E. coli</i> genome.
---

One or more products referenced in this document may be covered by a 3rd-party trademark.  
Please visit [www.neb.com/trademarks](http://www.neb.com/trademarks) for additional information.



Date 07 Oct 2022

---

Doreen Duquette  
Quality Approver

