

## New England Biolabs Certificate of Analysis

Product Name: NEB® 10-beta Competent *E. coli* (High Efficiency)  
 Catalog Number: C3019I  
 Lot Number: 10031508  
 Expiration Date: 11/2019  
 Storage Temperature: -80°C  
 Specification Version: PS-C3019H/I v1.0

NEB® 10-beta Competent <i>E. coli</i> (High Efficiency) Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
N3041AVIAL	pUC19 Vector	10025691	Pass
C3019IVIAL	NEB® 10-beta Competent <i>E. coli</i> (High Efficiency)	10017426	Pass
B9035SVIAL	NEB® 10-beta/Stable Outgrowth Medium	10014620	Pass

Assay Name/Specification	Lot # 10031508
<p><b>Antibiotic Sensitivity (Spectinomycin)</b>            15 µl of untransformed NEB® 10-beta Competent <i>E. coli</i> (High Efficiency) streaked onto a Rich Broth plate containing Spectinomycin will not form colonies after incubation for 16 hours at 37°C.</p>	Pass
<p><b>Antibiotic Sensitivity (Tetracycline)</b>            15 µl of untransformed NEB® 10-beta Competent <i>E. coli</i> (High Efficiency) streaked onto a Rich Broth plate containing Tetracycline will not form colonies after incubation for 16 hours at 37°C.</p>	Pass
<p><b>Blue-White Screening (α-complementation, Competent Cells)</b>            NEB® 10-beta Competent <i>E. coli</i> (High Efficiency) were shown to be suitable for blue/white screening by α-complementation of the β-galactosidase gene using pUC19.</p>	Pass
<p><b>Phage Resistance (φ 80)</b>            15 µl of untransformed NEB® 10-beta Competent <i>E. coli</i> (High Efficiency) streaked onto a Rich Broth plate does not support plaque formation by phage φ 80 after incubation for 16 hours at 37°C.</p>	Pass
<p><b>Transformation Efficiency</b>            50 µl of NEB® 10-beta Competent <i>E. coli</i> (High Efficiency) cells were transformed with 100 pg of pUC19 DNA using the transformation protocol provided. Incubation overnight on LB-Ampicillin plates at 37°C resulted in &gt;1 x 10e9 cfu/µg of DNA.</p>	Pass

Assay Name/Specification	Lot # 10031508
<p><b>Antibiotic Sensitivity (Kanamycin)</b> 15 µl of untransformed NEB® 10-beta Competent E. coli (High Efficiency) streaked onto a Rich Broth plate containing Kanamycin will not form colonies after incubation for 16 hours at 37°C.</p>	<b>Pass</b>
<p><b>Antibiotic Sensitivity (Nitrofurantoin)</b> 15 µl of untransformed NEB® 10-beta Competent E. coli (High Efficiency) streaked onto a Rich Broth plate containing Nitrofurantoin will not form colonies after incubation for 16 hours at 37°C.</p>	<b>Pass</b>
<p><b>Antibiotic Resistance (Streptomycin)</b> 15 µl of untransformed NEB® 10-beta Competent E. coli (High Efficiency) streaked onto a Rich Broth plate containing Streptomycin will form colonies after incubation for 16 hours at 37°C.</p>	<b>Pass</b>
<p><b>Antibiotic Sensitivity (Ampicillin)</b> 15 µl of untransformed NEB® 10-beta Competent E. coli (High Efficiency) streaked onto a Rich Broth plate containing Ampicillin will not form colonies after incubation for 16 hours at 37°C.</p>	<b>Pass</b>
<p><b>Antibiotic Sensitivity (Chloramphenicol)</b> 15 µl of untransformed NEB® 10-beta Competent E. coli (High Efficiency) streaked onto a Rich Broth plate containing Chloramphenicol will not form colonies after incubation for 16 hours at 37°C.</p>	<b>Pass</b>

This product has been tested and shown to be in compliance with all specifications.



Lixin An  
Production Scientist  
12 Oct 2018



Nick Privitera  
Packaging Quality Control Inspector  
06 Dec 2018