

New England Biolabs Certificate of Analysis

Product Name: *Taq 2X Master Mix*
Catalog Number: *M0270L*
Concentration: *2 X Concentrate*
Packaging Lot Number: *10151886*
Expiration Date: *01/2024*
Storage Temperature: *-20°C*
Specification Version: *PS-M0270S/L v2.0*
Composition (1X): *10 mM Tris-HCl (pH 8.6 @ 25°C), 50 mM KCl, 1.5 mM MgCl₂, 0.2 mM dATP, 0.2 mM dCTP, 0.2 mM dGTP, 0.2 mM dTTP, 5 % Glycerol, 0.08 % IGEPAL® CA-630, 0.05 % Tween® 20, 25 units/ml Taq DNA Polymerase*

Taq 2X Master Mix Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
M0270SVIAL	Taq 2X Master Mix	10132058	Pass
B9021SVIAL	Magnesium Chloride (MgCl ₂) Solution	10135558	Pass

Assay Name/Specification	Lot # 10151886
<p>qPCR DNA Contamination (E. coli Genomic) A minimum of 5 units of Taq DNA Polymerase is screened for the presence of E. coli genomic DNA using SYBR® Green qPCR with primers specific for the E. coli 16S rRNA locus. Results are quantified using a standard curve generated from purified E. coli genomic DNA. The measured level of E. coli genomic DNA contamination is ≤ 1 E. coli genome.</p>	Pass
<p>RNase Activity (Extended Digestion) A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 µl of Taq 2X Master Mix is incubated at 37°C. After incubation for 4 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.</p>	Pass
<p>Protein Purity Assay (SDS-PAGE) Taq DNA Polymerase is ≥ 99% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.</p>	Pass
<p>Phosphatase Activity (pNPP) A 200 µl reaction in 1M Diethanolamine, pH 9.8, 0.5 mM MgCl₂ containing 2.5 mM p-Nitrophenyl Phosphate (pNPP) and a minimum of 100 units of Taq DNA Polymerase</p>	Pass


Assay Name/Specification	Lot # 10151886
<p>incubated for 4 hours at 37°C yields <0.0001 unit of alkaline phosphatase activity as determined by spectrophotometric analysis.</p>	
<p>PCR Amplification (5 kb Lambda, Master Mix) A 25 µl reaction in 1X Taq Master Mix and 0.2 µM primers containing 5 ng Lambda DNA for 25 cycles of PCR amplification results in the expected 5 kb product.</p>	Pass
<p>Non-Specific DNase Activity (16 hour, Buffer) A 50 µl reaction in 1X Taq Master Mix containing 1 µg of T3 or T7 DNA in addition to a reaction containing Lambda-HindIII DNA incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.</p>	Pass
<p>Endonuclease Activity (Nicking) A 50 µl reaction in ThermoPol® Reaction Buffer containing 1 µg of supercoiled PhiX174 DNA and a minimum of 20 units of Taq DNA Polymerase incubated for 4 hours at 37°C and 75°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.</p>	Pass
<p>Single Stranded DNase Activity (FAM-Labeled Oligo) A 50 µl reaction in ThermoPol® Reaction Buffer containing a 10 nM solution of a fluorescent internal labeled oligonucleotide and a minimum of 25 units of Taq DNA Polymerase incubated for 30 minutes at 37°C and 75°C yields <10% degradation as determined by capillary electrophoresis.</p>	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.



Christie Vazquez
Production Scientist
27 May 2022



Michael Tonello
Packaging Quality Control Inspector
27 May 2022