

New England Biolabs Certificate of Analysis

Product Name: AMV Reverse Transcriptase
Catalog Number: M0277L
Concentration: 10,000 U/ml
Unit Definition: One unit is defined as the amount of enzyme required to incorporate 1 nmol of dTTP into an acid-insoluble form in 10 minutes at 37°C.
Packaging Lot Number: 10076892
Expiration Date: 01/2022
Storage Temperature: -20°C
Storage Conditions: 200 mM KPO₄, 2 mM DTT, 0.2 % Triton®X-100, 50 % Glycerol, (pH 7.2 @ 25°C)
Specification Version: PS-M0277S/L v2.0

AMV Reverse Transcriptase Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
M0277LVIAL	AMV Reverse Transcriptase	10072040	Pass
B0277AVIAL	AMV Reverse Transcriptase Reaction Buffer	10072362	Pass

Assay Name/Specification	Lot # 10076892
<p>RNase Activity Assay (4 Hour Digestion) A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 µl of AMV Reverse Transcriptase is incubated at 37°C. After incubation for 4 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.</p>	Pass
<p>Endonuclease Activity (Nicking) A 50 µl reaction in AMV Reverse Transcriptase Reaction Buffer containing 1 µg of supercoiled PhiX174 DNA and a minimum of 50 units of AMV Reverse Transcriptase incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.</p>	Pass
<p>Exonuclease Activity (Radioactivity Release) A 50 µl reaction in AMV Reverse Transcriptase Reaction Buffer containing 1 µg of a mixture of single and double-stranded [³H] E. coli DNA and a minimum of 50 units of AMV Reverse Transcriptase incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.</p>	Pass
<p>Non-Specific DNase Activity (16 Hour)</p>	Pass

Assay Name/Specification	Lot # 10076892
A 50 µl reaction in NEBuffer 2 containing 1 µg of T3 or T7 DNA in addition to a reaction containing Lambda-HindIII DNA and a minimum of 10 units of AMV Reverse Transcriptase incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.



Christie Vazquez
Production Scientist
22 Mar 2021



Michael Tonello
Packaging Quality Control Inspector
22 Mar 2021