

New England Biolabs Certificate of Analysis

Product Name: Multiplex PCR 5X Master Mix
Catalog Number: M0284S
Concentration: 5 X Concentrate
Packaging Lot Number: 10146119
Expiration Date: 03/2024
Storage Temperature: -20°C
Specification Version: PS-M0284S v2.0
Composition (1X): 20 mM Tris-HCl (pH 8.9 @ 25°C), 50 mM KCl, 30 mM NH₄Cl, 2.5 mM MgCl₂, 0.3 mM dATP, 0.3 mM dCTP, 0.3 mM dGTP, 0.3 mM dTTP, 3.2 % Glycerol, 0.08 % IGEPAL[®] CA-630, 0.07 % Tween[®] 20, 67 units/ml Taq DNA Polymerase

Multiplex PCR 5X Master Mix Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
M0284SVIAL	Multiplex PCR 5X Master Mix	10143628	Pass

Assay Name/Specification	Lot # 10146119
<p>Endonuclease Activity (Nicking) A 50 µl reaction in ThermoPol[®] Reaction Buffer containing 1 µg of supercoiled PhiX174 DNA and a minimum of 20 units of Taq DNA Polymerase incubated for 4 hours at 37°C and 75°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.</p>	Pass
<p>Single Stranded DNase Activity (FAM-Labeled Oligo) A 50 µl reaction in ThermoPol[®] Reaction Buffer containing a 10 nM solution of a fluorescent internal labeled oligonucleotide and a minimum of 25 units of Taq DNA Polymerase incubated for 30 minutes at 37°C and 75°C yields <10% degradation as determined by capillary electrophoresis.</p>	Pass
<p>Non-Specific DNase Activity (16 hour, Buffer) A 50 µl reaction in 2X Multiplex PCR Master Mix containing 1 µg of T3 or T7 DNA in addition to a reaction containing Lambda-HindIII DNA incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.</p>	Pass
<p>qPCR DNA Contamination (E. coli Genomic) A minimum of 5 units of Taq DNA Polymerase is screened for the presence of E. coli genomic DNA using SYBR[®] Green qPCR with primers specific for the E. coli 16S rRNA</p>	Pass


Assay Name/Specification	Lot # 10146119
<p>locus. Results are quantified using a standard curve generated from purified E. coli genomic DNA. The measured level of E. coli genomic DNA contamination is ≤ 1 E. coli genome.</p>	
<p>RNase Activity (Extended Digestion) A 10 μl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 μl of Multiplex PCR 5X Master Mix is incubated at 37°C. After incubation for 4 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.</p>	Pass
<p>Protein Purity Assay (SDS-PAGE) Taq DNA Polymerase is $\geq 99\%$ pure as determined by SDS-PAGE analysis using Coomassie Blue detection.</p>	Pass
<p>Phosphatase Activity (pNPP) A 200 μl reaction in 1M Diethanolamine, pH 9.8, 0.5 mM MgCl₂ containing 2.5 mM p-Nitrophenyl Phosphate (pNPP) and a minimum of 100 units of Taq DNA Polymerase incubated for 4 hours at 37°C yields <0.0001 unit of alkaline phosphatase activity as determined by spectrophotometric analysis.</p>	Pass
<p>PCR Amplification (15-plex PCR, Master Mix) A 25 μl reaction in 1X Multiplex PCR Master Mix and 0.15 μM primer mix containing 10 ng Human Genomic DNA for 35 cycles of PCR amplification results in the expected 15 products.</p>	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.



Christie Vazquez
Production Scientist
20 Apr 2022



Erin Varney
Packaging Quality Control Inspector
20 Apr 2022