

New England Biolabs Certificate of Analysis

Product Name: RNase Inhibitor, Murine
Catalog Number: M0314L
Concentration: 40,000 U/ml
Unit Definition: One unit is defined as the amount of Murine RNase Inhibitor required to inhibit the activity of 5ng of RNase A by 50%. Activity is measured by the inhibition of hydrolysis of cytidine 2', 3'-cyclic monophosphate by RNase A.
Lot Number: 10052579
Expiration Date: 08/2021
Storage Temperature: -20°C
Storage Conditions: 50 mM KCl, 20 mM HEPES (pH 7.6), 8 mM DTT, 50 % Glycerol
Specification Version: PS-M0314S/L v1.0


RNase Inhibitor, Murine Component List

| NEB Part Number | Component Description | Lot Number | Individual QC Result |
|-----------------|-------------------------|------------|----------------------|
| M0314LVIAL | RNase Inhibitor, Murine | 10052577 | Pass |

| Assay Name/Specification | Lot # 10052579 |
|--|----------------|
| Endonuclease Activity (Nicking) A 50 µl reaction in NEBuffer 4 containing 1 µg of supercoiled PhiX174 DNA and a minimum of 40 units of RNase Inhibitor, Murine incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis. | Pass |
| Exonuclease Activity (Radioactivity Release) A 50 µl reaction in NEBuffer 4 containing 1 µg of a mixture of single and double-stranded [³ H] E. coli DNA and a minimum of 200 units of RNase Inhibitor, Murine incubated for 4 hours at 37°C releases <0.1% of the total radioactivity. | Pass |
| Latent RNase Activity (Extended Digest) A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 40 units of heat inactivated RNase Inhibitor, Murine is incubated at 37°C. After incubation for 4 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection. | Pass |
| Protein Purity Assay (SDS-PAGE) RNase Inhibitor, Murine is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection. | Pass |

| Assay Name/Specification | Lot # 10052579 |
|--|--------------------|
| <p>RNase Activity (Extended Digestion) A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 40 units of RNase Inhibitor, Murine is incubated at 37°C. After incubation for 4 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.</p> | <p>Pass</p> |

This product has been tested and shown to be in compliance with all specifications.



Dongxian Yue
Production Scientist
26 Aug 2019



Jay Minichiello
Packaging Quality Control Inspector
13 Sep 2019