

## New England Biolabs Certificate of Analysis

**Product Name:** *RNase Inhibitor, Murine*  
**Catalog Number:** *M0314L*  
**Concentration:** *40,000 U/ml*  
**Unit Definition:** *One unit is defined as the amount of Murine RNase Inhibitor required to inhibit the activity of 5ng of RNase A by 50%. Activity is measured by the inhibition of hydrolysis of cytidine 2', 3'-cyclic monophosphate by RNase A.*  
**Packaging Lot Number:** *10144159*  
**Expiration Date:** *12/2023*  
**Storage Temperature:** *-20°C*  
**Storage Conditions:** *50 mM KCl, 20 mM HEPES (pH 7.6), 8 mM DTT, 50 % Glycerol*  
**Specification Version:** *PS-M0314S/L v1.0*

### RNase Inhibitor, Murine Component List


NEB Part Number	Component Description	Lot Number	Individual QC Result
M0314LVIAL	RNase Inhibitor, Murine	10130308	Pass

Assay Name/Specification	Lot # 10144159
<b>Protein Purity Assay (SDS-PAGE)</b> RNase Inhibitor, Murine is $\geq 95\%$ pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	Pass
<b>RNase Activity (Extended Digestion)</b> A 10 $\mu$ l reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 40 units of RNase Inhibitor, Murine is incubated at 37°C. After incubation for 4 hours, $>90\%$ of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.	Pass
<b>Exonuclease Activity (Radioactivity Release)</b> A 50 $\mu$ l reaction in NEBuffer 4 containing 1 $\mu$ g of a mixture of single and double-stranded [ <sup>3</sup> H] E. coli DNA and a minimum of 200 units of RNase Inhibitor, Murine incubated for 4 hours at 37°C releases $<0.1\%$ of the total radioactivity.	Pass
<b>Endonuclease Activity (Nicking)</b> A 50 $\mu$ l reaction in NEBuffer 4 containing 1 $\mu$ g of supercoiled PhiX174 DNA and a minimum of 40 units of RNase Inhibitor, Murine incubated for 4 hours at 37°C results in $<10\%$ conversion to the nicked form as determined by agarose gel electrophoresis.	Pass

Assay Name/Specification	Lot # 10144159
<p><b>Latent RNase Activity (Extended Digest)</b> A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 40 units of heat inactivated RNase Inhibitor, Murine is incubated at 37°C. After incubation for 4 hours, &gt;90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.</p>	<p><b>Pass</b></p>

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit [www.neb.com/trademarks](http://www.neb.com/trademarks) for additional information.



Bhairavi Jani  
Production Scientist  
10 Mar 2022



Michael Tonello  
Packaging Quality Control Inspector  
10 Mar 2022