

## New England Biolabs Certificate of Analysis

**Product Name:** OneTaq® Hot Start Quick-Load® 2X Master Mix with Standard Buffer  
**Catalog Number:** M0488S  
**Concentration:** 2 X Concentrate  
**Packaging Lot Number:** 10115508  
**Expiration Date:** 02/2023  
**Storage Temperature:** -20°C  
**Specification Version:** PS-M0488S/L v2.0  
**Composition (1X):** 20 mM Tris-HCl (pH 8.9 @ 25°C), 22 mM KCl, 22 mM NH<sub>4</sub>Cl, 1.8 mM MgCl<sub>2</sub>, 0.2 mM dATP, 0.2 mM dCTP, 0.2 mM dGTP, 0.2 mM dTTP, 5 % Glycerol, 0.06 % IGEPAL® CA-630, 0.05 % Tween® 20, 1 X Xylene cyanol, 1 X Tartrazine, 25 units/ml OneTaq® Hot Start DNA Polymerase

OneTaq® Hot Start Quick-Load® 2X Master Mix with Standard Buffer Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
M0488SVIAL	OneTaq® Hot Start Quick-Load® 2X Master Mix with Standard Buffer	10096420	Pass

Assay Name/Specification	Lot # 10115508
<p><b>PCR Amplification (5 kb Lambda, Master Mix)</b>            A 25 µl reaction in 1X OneTaq® Hot Start Quick-Load® Master Mix with Standard Buffer and 0.2 µM primers containing 5 ng Lambda DNA for 25 cycles of PCR amplification results in the expected 5 kb product.</p>	Pass
<p><b>PCR Amplification (Hot Start 2 kb Lambda DNA)</b>            A 25 µl reaction in OneTaq® Standard Reaction Buffer in the presence of 200 µM dNTPs and 0.2 µM primers containing 10 pg Lambda DNA and 50 ng Human Genomic DNA with 0.625 units of OneTaq® Hot Start DNA Polymerase for 30 cycles of PCR amplification results in an increase in yield of the 2 kb Lambda product and a decrease in non-specific genomic bands when compared to a non-hot start control reaction.</p>	Pass
<p><b>RNase Activity (Extended Digestion)</b>            A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 µl of OneTaq® Hot Start Quick-Load® 2X Master Mix with Standard Buffer is incubated at 37°C. After incubation for 4 hours, &gt;90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.</p>	Pass
<p><b>Inhibition of Primer Extension (Hot Start, Radioactivity Incorporation)</b>            A 50 µl primer extension assay in ThermoPol® Reaction Buffer in the presence of 200</p>	Pass


Assay Name/Specification	Lot # 10115508
<p><math>\mu</math>M dNTPs including [<sup>3</sup>H]-dTTP, containing 15 nM primed single-stranded M13mp18 with 2.5 units of OneTaq<sup>®</sup> Hot Start DNA Polymerase incubated for 16 hours at 25°C yields &gt;95% inhibition when compared to a non-hot start control reaction.</p> <p><b>Non-Specific DNase Activity (16 hour, Buffer)</b> A 50 <math>\mu</math>l reaction in 1X OneTaq<sup>®</sup> Hot Start Quick-Load<sup>®</sup> Master Mix with Standard Buffer containing 1 <math>\mu</math>g of T3 or T7 DNA in addition to a reaction containing Lambda-HindIII DNA incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.</p>	<p><b>Pass</b></p>

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit [www.neb.com/trademarks](http://www.neb.com/trademarks) for additional information.



Christie Vazquez  
Production Scientist  
20 Jul 2021



Michael Tonello  
Packaging Quality Control Inspector  
20 Jul 2021