

New England Biolabs Certificate of Analysis


Product Name: Quick CIP
Catalog Number: M0525L
Concentration: 5,000 U/ml
Unit Definition: One unit is defined as the amount of enzyme that hydrolyzes 1 μ mol of p-Nitrophenyl Phosphate, PNPP in a total reaction volume of 1 ml in 1 minute at 37°C.
Packaging Lot Number: 10057243
Expiration Date: 10/2021
Storage Temperature: -20°C
Storage Conditions: 25 mM Tris-HCl , 1 mM MgCl₂ , 0.1 mM ZnCl₂ , 50 % Glycerol, (pH 7.5 @ 25°C)
Specification Version: PS-M0525S/L v1.0

| Quick CIP Component List | | | |
|--------------------------|-----------------------|------------|----------------------|
| NEB Part Number | Component Description | Lot Number | Individual QC Result |
| M0525LVIAL | Quick CIP | 10055534 | Pass |
| B7204SVIAL | CutSmart® Buffer | 10053982 | Pass |

| Assay Name/Specification | Lot # 10057243 |
|--|----------------|
| Exonuclease Activity (Radioactivity Release) A 50 μ l reaction in CutSmart® Buffer containing 1 μ g of a mixture of single and double-stranded [³ H] E. coli DNA and a minimum of 50 units of Quick CIP incubated for 4 hours at 37°C releases <0.1% of the total radioactivity. | Pass |
| Endonuclease Activity (Nicking) A 50 μ l reaction in CutSmart® Buffer containing 1 μ g of supercoiled PhiX174 DNA and a minimum of 50 units of Quick CIP incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis. | Pass |
| RNase Activity (Extended Digestion) A 10 μ l reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 μ l of Quick CIP is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using gel electrophoresis using fluorescent detection. | Pass |
| Non-Specific DNase Activity (16 Hour) A 50 μ l reaction in NEBuffer 4 containing 1 μ g of PhiX174-HaeIII DNA and a minimum | Pass |

| Assay Name/Specification | Lot # 10057243 |
|--|----------------|
| of 50 units of Quick CIP incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis. | |

This product has been tested and shown to be in compliance with all specifications.



Doreen Duquette
Production Scientist
30 Sep 2019



Michael Tonello
Packaging Quality Control Inspector
07 Nov 2019