

New England Biolabs Certificate of Analysis

Product Name: β -N-Acetylglucosaminidase S
Catalog Number: P0744S
Concentration: 4,000 U/ml
Unit Definition: One unit is defined as the amount of enzyme required to cleave > 95% of the terminal, non-reducing β -N-Acetylglucosamine from 1 nmol GlcNAc β 1-4GlcNAc β 1-4GlcNAc-7-amino-4-methylcoumarin (AMC), in 1 hour at 37°C in a total reaction volume of 10 μ l.
Lot Number: 10009609
Expiration Date: 09/2019
Storage Temperature: 4°C
Storage Conditions: 50 mM NaCl, 20 mM Tris-HCl, 1 mM EDTA, (pH 7.5 @ 25°C)
Specification Version: PS-P0744S/L v1.0

β-N-Acetylglucosaminidase S Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
P0744SVIAL	β -N-Acetylglucosaminidase S	10018430	Pass
B1727SVIAL	10X GlycoBuffer 1	10011906	Pass

Assay Name/Specification	Lot # 10009609
Protein Purity Assay (SDS-PAGE) β -N-Acetylglucosaminidase S is \geq 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	Pass
Glycosidase Activity (α1-3 Galactosidase) A 10 μ l reaction in Glyco Buffer 1 containing 1 nM of fluorescently-labeled α -Galactosidase substrate (Gal α 1-3Gal β 1-4GlcNAc-AMC) and 16 units of β -N-Acetylglucosaminidase S incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	Pass
Glycosidase Activity (α1-3 Mannosidase) A 10 μ l reaction in Glyco Buffer 1 containing 1 nM of fluorescently-labeled α -Mannosidase substrate (Man α 1-3Man β 1-4GlcNAc-AMC) and 16 units of β -N-Acetylglucosaminidase S incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	Pass
Glycosidase Activity (α1-6 Galactosidase) A 10 μ l reaction in Glyco Buffer 1 containing 1 nM of fluorescently-labeled	Pass

Assay Name/Specification	Lot # 10009609
<p>α-Galactosidase substrate (Galα1-6Galα1-6Glcα1-2Fru-AMC) and 16 units of β-N-Acetylglucosaminidase S incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	
<p>Glycosidase Activity (α1-6 Mannosidase) A 10 μl reaction in Glyco Buffer 1 containing 1 nM of fluorescently-labeled α-Mannosidase substrate (Manα1-6Manα1-6(Manα1-3)Man-AMC) and 16 units of β-N-Acetylglucosaminidase S incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass
<p>Glycosidase Activity (α-N-Acetylgalactosaminidase) A 10 μl reaction in Glyco Buffer 1 containing 1 nM of fluorescently-labeled α-N-Acetylgalactosaminidase substrate (GalNAcα1-3(Fucα1-2)Galβ1-4Glc-AMC) and 16 units of β-N-Acetylglucosaminidase S incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass
<p>Glycosidase Activity (β-Mannosidase) A 10 μl reaction in Glyco Buffer 1 containing 1 nM of fluorescently-labeled β-Mannosidase substrate (Manβ1-4Manβ1-4Man-AMC) and 16 units of β-N-Acetylglucosaminidase S incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass
<p>Glycosidase Activity (β-Xylosidase) A 10 μl reaction in Glyco Buffer 1 containing 1 nM of fluorescently-labeled β-Xylosidase substrate (Xylβ1-4Xylβ1-4Xylβ1-4Xyl-AMC) and 16 units of β-N-Acetylglucosaminidase S incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass
<p>Glycosidase Activity (β1-3 Galactosidase) A 10 μl reaction in Glyco Buffer 1 containing 1 nM of fluorescently-labeled β-Galactosidase substrate (Galβ1-3GlcNAcβ1-4Galβ1-4Glc-AMC) and 16 units of β-N-Acetylglucosaminidase S incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass
<p>Glycosidase Activity (β1-4 Galactosidase) A 10 μl reaction in Glyco Buffer 1 containing 1 nM of fluorescently-labeled β-Galactosidase substrate (Galβ1-4GlcNAcβ1-3Galβ1-4Glc-AMC) and 16 units of β-N-Acetylglucosaminidase S incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass
<p>Glycosidase Activity (β-N-Acetylgalactosaminidase) A 10 μl reaction in Glyco Buffer 1 containing 1 nM of fluorescently-labeled β-N-Acetylgalactosaminidase substrate (GalNAcβ1-4Galβ1-4Glc-AMC) and 16 units of</p>	Pass

Assay Name/Specification	Lot # 10009609
<p>β-N-Acetylglucosaminidase S incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	
<p>Protease Activity (SDS-PAGE) A 20 μl reaction in 1X Glyco Buffer 1 containing 24 μg of a standard mixture of proteins and a minimum of 24 units of β-N-Acetylglucosaminidase S incubated for 20 hours at 37°C, results in no detectable degradation of the protein mixture as determined by SDS-PAGE with Coomassie Blue detection.</p>	Pass
<p>Glycosidase Activity (Endo F1, F2, H) A 10 μl reaction in Glyco Buffer 1 containing 1 nM of fluorescently-labeled Endo F1, F2, H substrate (Dansylated invertase high mannose) and 16 units of β-N-Acetylglucosaminidase S incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass
<p>Glycosidase Activity (Endo F2, F3) A 10 μl reaction in Glyco Buffer 1 containing 1 nM of fluorescently-labeled Endo F2, F3 substrate (Dansylated fibrinogen biantennary) and 16 units of β-N-Acetylglucosaminidase S incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass
<p>Glycosidase Activity (PNGase F) A 10 μl reaction in Glyco Buffer 1 containing 1 nM of fluorescently-labeled PNGase F substrate (Fluoresceinated fetuin triantennary) and 16 units of β-N-Acetylglucosaminidase S incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass
<p>Glycosidase Activity (α-Glucosidase) A 10 μl reaction in Glyco Buffer 1 containing 1 nM of fluorescently-labeled α-Glucosidase substrate (Glcα1-6Glcα1-4Glc-AMC) and 16 units of β-N-Acetylglucosaminidase S incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass
<p>Glycosidase Activity (α-Neuraminidase) A 10 μl reaction in Glyco Buffer 1 containing 1 nM of fluorescently-labeled α-Neuraminidase substrate (Neu5Acα2-3Galβ1-3GlcNAcβ1-3Galβ1-4Glc-AMC) and 16 units of β-N-Acetylglucosaminidase S incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass
<p>Glycosidase Activity (α1-2 Fucosidase) A 10 μl reaction in Glyco Buffer 1 containing 1 nM of fluorescently-labeled α-Fucosidase substrate (Fucα1-2Galβ1-4Glc-AMC) and 16 units of β-N-Acetylglucosaminidase S incubated for 20 hours at 37°C results in no detectable</p>	Pass

Assay Name/Specification	Lot # 10009609
<p>activity as determined by thin layer chromatography.</p> <p>Glycosidase Activity (α1-3 Fucosidase) A 10 μl reaction in Glyco Buffer 1 containing 1 nM of fluorescently-labeled α-Fucosidase substrate (Fuca1-3Galβ1-4GlcNAcβ1-3Galβ1-4Glc-AMC) and 16 units of β-N-Acetylglucosaminidase S incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	<p>Pass</p>

This product has been tested and shown to be in compliance with all specifications.



Alicia Bielik
Production Scientist
22 Jun 2018



Michael Tonello
Packaging Quality Control Inspector
09 Oct 2018