

## New England Biolabs Certificate of Analysis

**Product Name:** Bsp1286I  
**Catalog #:** R0120S/L  
**Concentration:** 5,000 units/ml  
**Unit Definition:** One unit is defined as the amount of enzyme required to digest 1 µg of Lambda DNA in 1 hour at 37°C in 50 µl of reaction buffer.  
**Lot #:** 0191503  
**Assay Date:** 03/2015  
**Expiration Date:** 3/2017  
**Storage Temp:** -20 °C  
**Specification Version:** PS-R0120S/L v1.0  
**Effective Date:** 12 Apr 2013

Assay Name/Specification (minimum release criteria)	Lot #0191503
<b>Exonuclease Activity (Radioactivity Release)</b> - A 50 µl reaction in CutSmart™ Buffer containing 1 µg of a mixture of single and double-stranded [ <sup>3</sup> H] <i>E. coli</i> DNA and a minimum of 50 units of Bsp1286I incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	<b>Pass</b>
<b>Ligation and Recutting (Terminal Integrity)</b> - After a 10-fold over-digestion of Lambda DNA with Bsp1286I, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with Bsp1286I.	<b>Pass</b>
<b>Non-Specific DNase Activity (16 Hour)</b> - A 50 µl reaction in CutSmart™ Buffer containing 1 µg of Lambda DNA and a minimum of 5 Units of Bsp1286I incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	<b>Pass</b>

\* The BSA in this product has been granted an EDQM "Certificate of Suitability" from the European Directorate for the Quality of Medicines (# R1-CEP-2003-204-Rev00) and has been granted a USDA Certificate for Export of Bovine Blood Plasma/Serum for Manufacture into Pharmaceutical Products.

M. W. Southworth



Authorized by  
Maurice Southworth  
12 Apr 2013

Inspected by  
Leo Formenoy  
06 Mar 2015

