

New England Biolabs Certificate of Analysis


Product Name: BbsI
Catalog Number: R0539L
Concentration: 10,000 U/ml
Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg of Lambda DNA in 1 hour at 37°C in a total reaction volume of 50 µl.
Lot Number: 10044977
Expiration Date: 05/2020
Storage Temperature: -80°C
Storage Conditions: 300 mM KCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 300 µg/ml BSA
Specification Version: PS-R0539S/L v2.0

| BbsI Component List | | | |
|---------------------|------------------------------|------------|----------------------|
| NEB Part Number | Component Description | Lot Number | Individual QC Result |
| R0539LVIAL | BbsI | 10044978 | Pass |
| B7202SVIAL | NEBuffer™ 2.1 | 10035939 | Pass |
| B7024SVIAL | Gel Loading Dye, Purple (6X) | 10038712 | Pass |

| Assay Name/Specification | Lot # 10044977 |
|---|----------------|
| <p>Endonuclease Activity (Nicking) A 50 µl reaction in NEBuffer 2.1 containing 1 µg of supercoiled pUC19 DNA and a minimum of 10 units of BbsI incubated for 4 hours at 37°C results in <20% conversion to the nicked form as determined by agarose gel electrophoresis.</p> | Pass |
| <p>Exonuclease Activity (Radioactivity Release) A 50 µl reaction in NEBuffer 2.1 containing 1 µg of a mixture of single and double-stranded [³H] E. coli DNA and a minimum of 50 units of BbsI incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.</p> | Pass |
| <p>Ligation and Recutting (Terminal Integrity) After a 20-fold over-digestion of Lambda DNA with BbsI, >95% of the DNA fragments can be ligated with T4 DNA ligase in 4 hours at 25°C. Of these ligated fragments, >95% can be recut with BbsI.</p> | Pass |
| <p>Non-Specific DNase Activity (16 Hour) A 50 ul reaction in NEBuffer 2.1 containing 1 ug of Lambda DNA and a minimum of 50 units of BbsI incubated for 16 hours at 37°C results in a DNA pattern free of</p> | Pass |

| Assay Name/Specification | Lot # 10044977 |
|---|----------------|
| detectable nuclease degradation as determined by agarose gel electrophoresis. | |

This product has been tested and shown to be in compliance with all specifications.



Doreen Duquette
Production Scientist
06 Feb 2019



Jay Minichiello
Packaging Quality Control Inspector
14 Jun 2019