

New England Biolabs Certificate of Analysis

Product Name: PmeI
Catalog Number: R0560S
Concentration: 10,000 U/ml
Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg of Lambda DNA in 1 hour at 37°C in a total reaction volume of 50 µl.
Packaging Lot Number: 10061234
Expiration Date: 12/2021
Storage Temperature: -20°C
Storage Conditions: 100 mM NaCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 200 µg/ml BSA
Specification Version: PS-R0560S/L v1.0

PmeI Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
R0560SVIAL	PmeI	10060897	Pass
B7204SVIAL	CutSmart® Buffer	10055736	Pass
B7024SVIAL	Gel Loading Dye, Purple (6X)	10055732	Pass


Assay Name/Specification	Lot # 10061234
Blue-White Screening (Terminal Integrity) A sample of pNEB193 vector linearized with a 10-fold excess of PmeI, religated and transformed into an E. coli strain expressing the LacZ beta fragment gene results in <1% white colonies.	Pass
Endonuclease Activity (Nicking) A 50 µl reaction in CutSmart™ Buffer containing 1 µg of supercoiled PhiX174 DNA and a minimum of 10 Units of PmeI incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
Exonuclease Activity (Radioactivity Release) A 50 µl reaction in CutSmart™ Buffer containing 1 µg of a mixture of single and double-stranded [³ H] E. coli DNA and a minimum of 100 units of PmeI incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass
Non-Specific DNase Activity (16 Hour) A 50 µl reaction in CutSmart™ Buffer containing 1 µg of Lambda DNA and a minimum of 10 Units of PmeI incubated for 16 hours at 37°C results in a DNA pattern free of	Pass

Assay Name/Specification	Lot # 10061234
<p>detectable nuclease degradation as determined by agarose gel electrophoresis.</p> <p>Ligation and Recutting (Terminal Integrity) After a 10-fold over-digestion of Lambda DNA with PmeI, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with PmeI.</p>	<p>Pass</p>

This product has been tested and shown to be in compliance with all specifications.



Jianying Luo
Production Scientist
27 Nov 2019



Michael Tonello
Packaging Quality Control Inspector
30 Dec 2019