

New England Biolabs Certificate of Analysis

Product Name: Cac8I
Catalog #: R0579S/L
Concentration: 5,000 units/ml
Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg of Lambda DNA in 1 hour at 37°C in a total reaction volume of 50 µl.
Lot #: 0271803
Assay Date: 03/2018
Expiration Date: 3/2020
Storage Temp: -20°C
Storage Conditions: 150 mM KCl , 10 mM Tris-HCl (7.4), 1 mM DTT , 0.1 mM EDTA , 50 % Glycerol , 0.10 % TritonX-100
Specification Version: PS-R0579S/L v3.0
Effective Date: 01 Dec 2016

Assay Name/Specification (minimum release criteria)	Lot #0271803
Exonuclease Activity (Radioactivity Release) - A 50 µl reaction in CutSmart® Buffer containing 1 µg of a mixture of single and double-stranded [³ H] <i>E. coli</i> DNA and a minimum of 15 units of Cac8I incubated for 4 hours at 37°C releases <0.2% of the total radioactivity.	Pass
Ligation and Recutting (Terminal Integrity) - After a 5-fold over-digestion of Lambda DNA with Cac8I, ~75% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with Cac8I.	Pass
Non-Specific DNase Activity (16 Hour) - A 50 µl reaction in CutSmart® Buffer containing 1 µg of Lambda DNA and a minimum of 25 units of Cac8I incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass



Authorized by
Derek Robinson
01 Dec 2016



Inspected by
Stephanie Cornelio
07 Mar 2018

