

## New England Biolabs Certificate of Analysis

**Product Name:** Sall-HF®  
**Catalog Number:** R3138S  
**Concentration:** 20,000 U/ml  
**Unit Definition:** One unit is defined as the amount of enzyme required to digest 1 µg of Lambda DNA (HindIII digest) in 1 hour at 37°C in a total reaction volume of 50 µl.  
**Lot Number:** 10034666  
**Expiration Date:** 12/2020  
**Storage Temperature:** -20°C  
**Storage Conditions:** 50 mM KCl, 10 mM Tris-HCl (pH 7.5), 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 300 µg/ml BSA  
**Specification Version:** PS-R3138S/L v1.0

| Sall-HF® Component List |                              |            |                      |
|-------------------------|------------------------------|------------|----------------------|
| NEB Part Number         | Component Description        | Lot Number | Individual QC Result |
| R3138SVIAL              | Sall-HF®                     | 10030763   | Pass                 |
| B7204SVIAL              | CutSmart® Buffer             | 10031565   | Pass                 |
| B7024SVIAL              | Gel Loading Dye, Purple (6X) | 10021138   | Pass                 |

| Assay Name/Specification  | Lot # 10034666 |
|---|----------------|
| <b>Ligation and Recutting (Terminal Integrity)</b><br>After a 50-fold over-digestion of Adenovirus-2 DNA with Sall-HF™, >95% of the DNA fragments can be ligated with T4 DNA ligase in 4 hours at 25°C. Of these ligated fragments, >95% can be recut with Sall-HF™.                                    | Pass           |
| <b>Non-Specific DNase Activity (16 Hour)</b><br>A 50 µl reaction in CutSmart™ Buffer containing 1 µg of pBR322 DNA and a minimum of 200 Units of Sall-HF™ incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis. | Pass           |
| <b>Exonuclease Activity (Radioactivity Release)</b><br>A 50 µl reaction in CutSmart™ Buffer containing 1 µg of a mixture of single and double-stranded [ <sup>3</sup> H] E. coli DNA and a minimum of 200 units of Sall-HF™ incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.    | Pass           |
| <b>Blue-White Screening (Terminal Integrity)</b><br>A sample of pUC19 vector linearized with a 10-fold excess of Sall-HF™, religated  | Pass           |

| Assay Name/Specification  | Lot # 10034666     |
|---|--------------------|
| <p>and transformed into an E. coli strain expressing the LacZ beta fragment gene results in &lt;1% white colonies.</p> <p><b>Endonuclease Activity (Nicking)</b><br/>A 50 µl reaction in CutSmart™ Buffer containing 1 µg of supercoiled PhiX174 DNA and a minimum of 20 Units of Sall-HF™ incubated for 4 hours at 37°C results in &lt;10% conversion to the nicked form as determined by agarose gel electrophoresis.</p> | <p><b>Pass</b></p> |

This product has been tested and shown to be in compliance with all specifications.



Jianying Luo  
Production Scientist  
05 Dec 2018



Michael Tonello  
Packaging Quality Control Inspector  
30 Jan 2019