

## New England Biolabs Certificate of Analysis

**Product Name:** EagI-HF<sup>®</sup>  
**Catalog Number:** R3505M  
**Concentration:** 100,000 U/ml  
**Unit Definition:** One unit is defined as the amount of enzyme required to digest 1 µg of pXba DNA in 1 hour at 37°C in a total reaction volume of 50 µl.  
**Packaging Lot Number:** 10151155  
**Expiration Date:** 02/2024  
**Storage Temperature:** -80°C  
**Storage Conditions:** 500 mM NaCl, 10 mM Tris-HCl, 1 mM DTT, 0.1 mM EDTA, 50 % Glycerol, 200 µg/ml BSA, (pH 7.4 @ 25°C)  
**Specification Version:** PS-R3505M v3.0

| EagI-HF <sup>®</sup> Component List |                               |            |                      |
|-------------------------------------|-------------------------------|------------|----------------------|
| NEB Part Number                     | Component Description         | Lot Number | Individual QC Result |
| R3505MVIAL                          | EagI-HF <sup>®</sup>          | 10139929   | Pass                 |
| B7024AVIAL                          | Gel Loading Dye, Purple (6X)  | 10132772   | Pass                 |
| B6004SVIAL                          | rCutSmart <sup>™</sup> Buffer | 10146828   | Pass                 |

| Assay Name/Specification   | Lot # 10151155 |
|--|----------------|
| <b>Blue-White Screening (Terminal Integrity)</b><br>A sample of Litmus38i vector linearized with a 10-fold excess of EagI-HF <sup>™</sup> , religated and transformed into an E. coli strain expressing the LacZ beta fragment gene results in <1% white colonies.   | Pass           |
| <b>Ligation and Recutting (Terminal Integrity)</b><br>After a 20-fold over-digestion of pXba DNA with EagI-HF <sup>™</sup> , >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with EagI-HF <sup>™</sup> .                                      | Pass           |
| <b>Exonuclease Activity (Radioactivity Release)</b><br>A 50 µl reaction in CutSmart <sup>™</sup> Buffer containing 1 µg of a mixture of single and double-stranded [ <sup>3</sup> H] E. coli DNA and a minimum of 100 units of EagI-HF <sup>™</sup> incubated for 4 hours at 37°C releases <0.1% of the total radioactivity. | Pass           |
| <b>Non-Specific DNase Activity (16 Hour)</b><br>A 50 µl reaction in CutSmart <sup>™</sup> Buffer containing 1 µg of pXba DNA and a minimum of 100 Units of EagI-HF <sup>™</sup> incubated for 16 hours at 37°C results in a DNA pattern free   | Pass           |

| Assay Name/Specification   | Lot # 10151155 |
|--|----------------|
| of detectable nuclease degradation as determined by agarose gel electrophoresis.   |                |
| <p><b>Endonuclease Activity (Nicking)</b><br/>A 50 µl reaction in CutSmart™ Buffer containing 1 µg of supercoiled PhiX174 DNA and a minimum of 20 Units of EagI-HF™ incubated for 4 hours at 37°C results in &lt;20% conversion to the nicked form as determined by agarose gel electrophoresis.</p> | <b>Pass</b>    |
| <p><b>Protein Purity Assay (SDS-PAGE)</b><br/>EagI-HF™ is &gt;95% pure as determined by SDS PAGE analysis using Coomassie Blue detection.</p>  | <b>Pass</b>    |

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit [www.neb.com/trademarks](http://www.neb.com/trademarks) for additional information.



Penghua Zhang  
Production Scientist  
11 May 2022



Erin Varney  
Packaging Quality Control Inspector  
11 May 2022