

New England Biolabs Certificate of Analysis

Product Name: BsrGI-HF[®]
Catalog Number: R3575S
Concentration: 20,000 U/ml
Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg of Lambda DNA in 1 hour at 37°C in a total reaction volume of 50 µl.
Lot Number: 10052846
Expiration Date: 04/2021
Storage Temperature: -20°C
Storage Conditions: 50 mM NaCl , 10 mM Tris-HCl (pH 7.4), 1 mM DTT , 0.1 mM EDTA , 50 % Glycerol , 200 µg/ml BSA
Specification Version: PS-R3575S/L v1.0

BsrGI-HF [®] Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
R3575SVIAL	BsrGI-HF [®]	10043043	Pass
B7204SVIAL	CutSmart [®] Buffer	10046089	Pass
B7024SVIAL	Gel Loading Dye, Purple (6X)	10043911	Pass

Assay Name/Specification	Lot # 10052846
Endonuclease Activity (Nicking) A 50 µl reaction in CutSmart [™] Buffer containing 1 µg of supercoiled φX174 DNA and a minimum of 60 units of BsrGI-HF incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
Exonuclease Activity (Radioactivity Release) A 50 µl reaction in CutSmart [™] Buffer containing 1 µg of a mixture of single and double-stranded [³ H] E. coli DNA and a minimum of 100 units of BsrGI-HF incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass
Functional Test (15 minute Digest) A 50 µl reaction in CutSmart [™] Buffer containing 1 µg of Lambda DNA and 1 µl of BsrGI-HF incubated for 15 minutes at 37°C results in complete digestion as determined by agarose gel electrophoresis.	Pass
Ligation and Recutting (Terminal Integrity) After a 10-fold over-digestion of Lambda DNA with BsrGI-HF, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated	Pass

Assay Name/Specification	Lot # 10052846
fragments, >95% can be recut with BsrGI-HF.	
<p>Non-Specific DNase Activity (16 Hour) A 50 µl reaction in CutSmart™ Buffer containing 1 µg of Lambda DNA and a minimum of 100 units of BsrGI-HF incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.</p>	Pass
<p>Protein Purity Assay (SDS-PAGE) BsrGI-HF is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.</p>	Pass

This product has been tested and shown to be in compliance with all specifications.



Stephanie Cornelio
Production Scientist
23 Apr 2019



Jay Minichiello
Packaging Quality Control Inspector
16 Sep 2019