

**T4 Polynucleotide
Kinase Reaction Buffer**



1-800-632-7799
info@neb.com
www.neb.com



B0201S 001160719071

B0201S

4.0 ml T4 Polynucleotide Kinase Exp: 7/19

Reaction Buffer (10X) Lot: 0011607

Store at -20°C

Description: New England Biolabs supplies a 10X reaction buffer with all of its enzymes. At a 1X concentration this reaction buffer assures optimal activity of the enzyme.

1X T4 Polynucleotide Kinase Reaction Buffer:

70 mM Tris-HCl

10 mM MgCl₂

5 mM DTT

pH 7.6 @ 25°C

Supplied with enzyme as a 10X concentrated stock

Quality Control Assays

16-Hour Incubation: A 50 µl reaction containing this reaction buffer at a 1X concentration and 1 µg of HaeIII digested φX174 RF I DNA incubated for 16 hours resulted in no detectable non-specific nuclease degradation.

Endonuclease Activity: Incubation of this reaction buffer at a 1X concentration with 1 µg φX174 RF I DNA for 4 hours at 37°C in 50 µl reactions resulted in less than 5% conversion to RF II.



NEW ENGLAND BIOLABS® is a registered trademark of New England Biolabs, Inc.

This product is intended for research purposes only. This product is not intended to be used for therapeutic or diagnostic purposes in humans or animals.

CERTIFICATE OF ANALYSIS

**T4 Polynucleotide
Kinase Reaction Buffer**



1-800-632-7799
info@neb.com
www.neb.com



B0201S 001160719071

B0201S

4.0 ml T4 Polynucleotide Kinase Exp: 7/19

Reaction Buffer (10X) Lot: 0011607

Store at -20°C

Description: New England Biolabs supplies a 10X reaction buffer with all of its enzymes. At a 1X concentration this reaction buffer assures optimal activity of the enzyme.

1X T4 Polynucleotide Kinase Reaction Buffer:

70 mM Tris-HCl

10 mM MgCl₂

5 mM DTT

pH 7.6 @ 25°C

Supplied with enzyme as a 10X concentrated stock

Quality Control Assays

16-Hour Incubation: A 50 µl reaction containing this reaction buffer at a 1X concentration and 1 µg of HaeIII digested φX174 RF I DNA incubated for 16 hours resulted in no detectable non-specific nuclease degradation.

Endonuclease Activity: Incubation of this reaction buffer at a 1X concentration with 1 µg φX174 RF I DNA for 4 hours at 37°C in 50 µl reactions resulted in less than 5% conversion to RF II.



NEW ENGLAND BIOLABS® is a registered trademark of New England Biolabs, Inc.

This product is intended for research purposes only. This product is not intended to be used for therapeutic or diagnostic purposes in humans or animals.

CERTIFICATE OF ANALYSIS