

Taq DNA Ligase Reaction Buffer



1-800-632-7799
info@neb.com
www.neb.com



B0208S 002140117011

B0208S

6.0 ml *Taq* DNA Ligase Reaction Buffer (10X)

Store at -20°C Lot: 0021401 Exp: 1/17

Description: New England Biolabs supplies a 10X reaction buffer with all of its enzymes. At a 1X concentration this reaction buffer assures optimal activity of the enzyme.

1X *Taq* DNA Ligase Reaction Buffer:

20 mM Tris-HCl
25 mM potassium acetate
10 mM magnesium acetate
10 mM DTT
1 mM NAD
0.1% Triton X-100
pH 7.6 @ 25°C

Taq DNA Ligase Reaction Buffer



1-800-632-7799
info@neb.com
www.neb.com



B0208S 002140117011

B0208S

6.0 ml *Taq* DNA Ligase Reaction Buffer (10X)

Store at -20°C Lot: 0021401 Exp: 1/17

Description: New England Biolabs supplies a 10X reaction buffer with all of its enzymes. At a 1X concentration this reaction buffer assures optimal activity of the enzyme.

1X *Taq* DNA Ligase Reaction Buffer:

20 mM Tris-HCl
25 mM potassium acetate
10 mM magnesium acetate
10 mM DTT
1 mM NAD
0.1% Triton X-100
pH 7.6 @ 25°C

Quality Control Assays

16-Hour Incubation: A 50 μl reaction containing this reaction buffer at a 1X concentration and 1 μg of HaeIII digested ϕX174 RF I DNA incubated for 16 hours resulted in no detectable non-specific nuclease degradation.

Endonuclease Activity: Incubation of this reaction buffer at a 1X concentration with 1 μg ϕX174 RF I DNA for 4 hours at 37°C in 50 μl reactions resulted in less than 5% conversion to RF II.

CERTIFICATE OF ANALYSIS

Quality Control Assays

16-Hour Incubation: A 50 μl reaction containing this reaction buffer at a 1X concentration and 1 μg of HaeIII digested ϕX174 RF I DNA incubated for 16 hours resulted in no detectable non-specific nuclease degradation.

Endonuclease Activity: Incubation of this reaction buffer at a 1X concentration with 1 μg ϕX174 RF I DNA for 4 hours at 37°C in 50 μl reactions resulted in less than 5% conversion to RF II.

CERTIFICATE OF ANALYSIS