

# Exonuclease I Reaction Buffer



1-800-632-7799  
info@neb.com  
www.neb.com



B0293S 001160619061

## B0293S

6.0 ml Exonuclease I Reaction Buffer (10X)

Store at  $-20^{\circ}\text{C}$  Lot: 0011606 Exp: 6/19

**Description:** New England Biolabs supplies a 10X reaction buffer with all of its enzymes. At a 1X concentration this reaction buffer assures optimal activity of the enzyme.

### 1X Exonuclease I Reaction Buffer:

67 mM Glycine-KOH  
6.7 mM  $\text{MgCl}_2$   
10 mM 2-mercaptoethanol  
(pH 9.5 @  $25^{\circ}\text{C}$ )

### Quality Control Assay

**16-Hour Incubation:** A 50  $\mu\text{l}$  reaction containing this reaction buffer at a 1X concentration and 1  $\mu\text{g}$  of HaeIII digested  $\phi\text{X174}$  RF I DNA incubated for 16 hours resulted in no detectable non-specific nuclease degradation.

**Endonuclease Activity:** Incubation of this reaction buffer at a 1X concentration with 1  $\mu\text{g}$   $\phi\text{X174}$  RF I DNA for 4 hours at  $37^{\circ}\text{C}$  in 50  $\mu\text{l}$  reactions resulted in less than 5% conversion to RF II.



NEW ENGLAND BIOLABS® is a registered trademark of New England Biolabs, Inc.

This product is intended for research purposes only. This product is not intended to be used for therapeutic or diagnostic purposes in humans or animals.

CERTIFICATE OF ANALYSIS

# Exonuclease I Reaction Buffer



1-800-632-7799  
info@neb.com  
www.neb.com



B0293S 001160619061

## B0293S

6.0 ml Exonuclease I Reaction Buffer (10X)

Store at  $-20^{\circ}\text{C}$  Lot: 0011606 Exp: 6/19

**Description:** New England Biolabs supplies a 10X reaction buffer with all of its enzymes. At a 1X concentration this reaction buffer assures optimal activity of the enzyme.

### 1X Exonuclease I Reaction Buffer:

67 mM Glycine-KOH  
6.7 mM  $\text{MgCl}_2$   
10 mM 2-mercaptoethanol  
(pH 9.5 @  $25^{\circ}\text{C}$ )

### Quality Control Assay

**16-Hour Incubation:** A 50  $\mu\text{l}$  reaction containing this reaction buffer at a 1X concentration and 1  $\mu\text{g}$  of HaeIII digested  $\phi\text{X174}$  RF I DNA incubated for 16 hours resulted in no detectable non-specific nuclease degradation.

**Endonuclease Activity:** Incubation of this reaction buffer at a 1X concentration with 1  $\mu\text{g}$   $\phi\text{X174}$  RF I DNA for 4 hours at  $37^{\circ}\text{C}$  in 50  $\mu\text{l}$  reactions resulted in less than 5% conversion to RF II.



NEW ENGLAND BIOLABS® is a registered trademark of New England Biolabs, Inc.

This product is intended for research purposes only. This product is not intended to be used for therapeutic or diagnostic purposes in humans or animals.

CERTIFICATE OF ANALYSIS