

NEBNext®
Quick Ligation
Reaction Buffer



1-800-632-7799
info@neb.com
www.neb.com



B6058S 017170319031

B6058S

2 x 1.0 ml (2 ml) Lot: 0171703
Store at -20°C Exp: 3/19

Description: New England Biolabs supplies a 5X reaction buffer for use with Quick T4 DNA Ligase. At a 1X concentration this reaction buffer assures optimal activity of the enzyme.

1X NEBNext Quick Ligation Reaction Buffer:

66 mM Tris-HCl
10 mM MgCl₂
1 mM DTT
1 mM ATP
6% Polyethylene glycol (PEG 6000)
pH 7.6 @ 25°C

NEBNext®
Quick Ligation
Reaction Buffer



1-800-632-7799
info@neb.com
www.neb.com



B6058S 017170319031

B6058S

2 x 1.0 ml (2 ml) Lot: 0171703
Store at -20°C Exp: 3/19

Description: New England Biolabs supplies a 5X reaction buffer for use with Quick T4 DNA Ligase. At a 1X concentration this reaction buffer assures optimal activity of the enzyme.

1X NEBNext Quick Ligation Reaction Buffer:

66 mM Tris-HCl
10 mM MgCl₂
1 mM DTT
1 mM ATP
6% Polyethylene glycol (PEG 6000)
pH 7.6 @ 25°C

Quality Control Assay

16-Hour Incubation: 50 µl reactions containing this reaction buffer at 1X concentration and 1 µg of HindIII digested Lambda DNA incubated for 16 hours at 37°C results in no detectable non-specific nuclease degradation as determined by agarose gel electrophoresis. 50 µl reactions containing this reaction buffer at 1X concentration and 1 µg T3 DNA incubated for 16 hours at 37°C also results in no detectable non-specific nuclease degradation as determined by agarose gel electrophoresis.

Endonuclease Activity: Incubation of this reaction buffer at a 1X concentration with 1 µg of φX174 RF I DNA for 4 hours at 37°C in 50 µl reactions results in less than 10% conversion to RF II as determined by agarose gel electrophoresis.

RNase Activity: Incubation of this reaction buffer at 1X concentration with 40 ng of a FAM-labeled RNA transcript for 16 hours at 37°C results in no detectable RNase activity as determined by polyacrylamide gel electrophoresis.

Quality Control Assay

16-Hour Incubation: 50 µl reactions containing this reaction buffer at 1X concentration and 1 µg of HindIII digested Lambda DNA incubated for 16 hours at 37°C results in no detectable non-specific nuclease degradation as determined by agarose gel electrophoresis. 50 µl reactions containing this reaction buffer at 1X concentration and 1 µg T3 DNA incubated for 16 hours at 37°C also results in no detectable non-specific nuclease degradation as determined by agarose gel electrophoresis.

Endonuclease Activity: Incubation of this reaction buffer at a 1X concentration with 1 µg of φX174 RF I DNA for 4 hours at 37°C in 50 µl reactions results in less than 10% conversion to RF II as determined by agarose gel electrophoresis.

RNase Activity: Incubation of this reaction buffer at 1X concentration with 40 ng of a FAM-labeled RNA transcript for 16 hours at 37°C results in no detectable RNase activity as determined by polyacrylamide gel electrophoresis.

Phosphatase Activity: Incubation of this reaction buffer at a 1X concentration in protein phosphatase assay buffer (1 M diethanolamine @ pH 9.8 and 0.5 mM MgCl₂) containing 2.5 mM *p*-nitrophenyl phosphate at 37°C for 4 hours yields no detectable *p*-nitrophenylene anion as determined by spectrophotometric analysis at 405 nm.



NEW ENGLAND BIOLABS® and NEBNEXT® are registered trademarks of New England Biolabs, Inc.

This product is intended for research purposes only. This product is not intended to be used for therapeutic or diagnostic purposes in humans or animals.

CERTIFICATE OF ANALYSIS

Phosphatase Activity: Incubation of this reaction buffer at a 1X concentration in protein phosphatase assay buffer (1 M diethanolamine @ pH 9.8 and 0.5 mM MgCl₂) containing 2.5 mM *p*-nitrophenyl phosphate at 37°C for 4 hours yields no detectable *p*-nitrophenylene anion as determined by spectrophotometric analysis at 405 nm.



NEW ENGLAND BIOLABS® and NEBNEXT® are registered trademarks of New England Biolabs, Inc.

This product is intended for research purposes only. This product is not intended to be used for therapeutic or diagnostic purposes in humans or animals.

CERTIFICATE OF ANALYSIS