

CutSmart™ Buffer



1-800-632-7799
info@neb.com
www.neb.com



B7204S 005131116111

B7204S

5.0 ml

Lot: 0051311

Store at -20°C


Exp: 11/16

Description: New England Biolabs provides a color-coded 10X NEBuffer with each restriction endonuclease to ensure optimal (100%) activity. Most of our enzymes are supplied with one of four standard NEBuffers. Occasionally, an enzyme has specific buffer requirements not met by one of the four standard NEBuffers, in which case the enzyme is supplied with its own unique NEBuffer.

CutSmart™ Buffer



1-800-632-7799
info@neb.com
www.neb.com



B7204S 005131116111

B7204S

5.0 ml

Lot: 0051311

Store at -20°C

Exp: 11/16

Description: New England Biolabs provides a color-coded 10X NEBuffer with each restriction endonuclease to ensure optimal (100%) activity. Most of our enzymes are supplied with one of four standard NEBuffers. Occasionally, an enzyme has specific buffer requirements not met by one of the four standard NEBuffers, in which case the enzyme is supplied with its own unique NEBuffer.

1X CutSmart Buffer:

50 mM Potassium acetate
20 mM Tris-acetate
10 mM Magnesium acetate
100 µg/ml BSA
pH 7.9 @ 25°C
Supplied as a 10X concentrated stock

Quality Control

pH range: The pH of 10X CutSmart Buffer is between pH 7.8 and 8.0.

16-hour Non-specific Nuclease Activity Assay:

A 50 µl reaction in 1X CutSmart Buffer containing 1 µg of ϕ X HaeIII digested DNA incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.

Endonuclease (nicking) Activity Assay: A 50 µl reaction in 1X CutSmart Buffer containing 1 µg of supercoiled ϕ X174 DNA incubated for 4 hours at 37°C results in < 10% conversion to the nicked form as determined by agarose gel electrophoresis.

1X CutSmart Buffer:

50 mM Potassium acetate
20 mM Tris-acetate
10 mM Magnesium acetate
100 µg/ml BSA
pH 7.9 @ 25°C
Supplied as a 10X concentrated stock

Quality Control

pH range: The pH of 10X CutSmart Buffer is between pH 7.8 and 8.0.

16-hour Non-specific Nuclease Activity Assay:

A 50 µl reaction in 1X CutSmart Buffer containing 1 µg of ϕ X HaeIII digested DNA incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.

Endonuclease (nicking) Activity Assay: A 50 µl reaction in 1X CutSmart Buffer containing 1 µg of supercoiled ϕ X174 DNA incubated for 4 hours at 37°C results in < 10% conversion to the nicked form as determined by agarose gel electrophoresis.

Buffer Functional Assay: A 50 µl reaction in 1X CutSmart Buffer containing 1 µg of λ DNA and 1 unit of MscI, or 1 µg of λ DNA (*N*⁶-methyladenine-free) and 1 unit of ClaI, incubated for 1 hour at 37°C results in complete digestion of the substrate DNA as determined by agarose gel electrophoresis.

RNase Activity (Extended Digestion): A 10 µl reaction in 1X CutSmart Buffer with 40 ng RNA transcript is incubated for 16 hours at 37°C. After incubation for 16 hours, no detectable degradation of the RNA is observed as determined by gel electrophoresis using fluorescent detection.



NEW ENGLAND BIOLABS® is a registered trademark of New England Biolabs, Inc.
CUTSMART™ is a trademark of New England Biolabs, Inc.

Buffer Functional Assay: A 50 µl reaction in 1X CutSmart Buffer containing 1 µg of λ DNA and 1 unit of MscI, or 1 µg of λ DNA (*N*⁶-methyladenine-free) and 1 unit of ClaI, incubated for 1 hour at 37°C results in complete digestion of the substrate DNA as determined by agarose gel electrophoresis.

RNase Activity (Extended Digestion): A 10 µl reaction in 1X CutSmart Buffer with 40 ng RNA transcript is incubated for 16 hours at 37°C. After incubation for 16 hours, no detectable degradation of the RNA is observed as determined by gel electrophoresis using fluorescent detection.



NEW ENGLAND BIOLABS® is a registered trademark of New England Biolabs, Inc.
CUTSMART™ is a trademark of New England Biolabs, Inc.

CERTIFICATE OF ANALYSIS

CERTIFICATE OF ANALYSIS