

BSA,  
Molecular Biology  
Grade



1-800-632-7799  
info@neb.com  
www.neb.com



B9000S 006150618061

**B9000S**

**12 mg BSA, Molecular Biology Grade Exp: 6/18**  
**20 mg/ml Lot: 0061506 Store at -20°C**

**Description:** Bovine Serum Albumin (BSA) is supplied with some products to prevent adhesion of the enzyme to reaction tubes and pipette surfaces. BSA also stabilizes some proteins during incubation.

Supplied in: 20 mM Tris-HCl, 100 mM KCl, 0.1 mM EDTA and 50% glycerol (pH 8.0 @ 25°C).

**Quality Controls Assays**

**Protein Concentration ( $A_{280}$ ):** The concentration of BSA is 20 mg/ml +/- 5% as determined by UV absorption at 280 nm. Protein concentration is determined by the Pace method using the extinction coefficient of 42,925 and molecular weight of 66,464 daltons for BSA (1).

**Non-Specific DNase Activity (16 hour):** A 50  $\mu$ l reaction in NEBuffer 4 containing 1  $\mu$ g of Lambda DNA (HindIII digest) and a minimum of 100  $\mu$ g of BSA incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.

**RNase Activity (2 Hour Digestion):** A 10  $\mu$ l reaction in NEBuffer 4 containing 40 ng of fluorescein labeled RNA transcript and a minimum of 20  $\mu$ g of BSA incubated for 2 hours at 37°C results in no detectable degradation of the RNA as determined by gel electrophoresis using fluorescence detection.

**Quality Controls Assays**

**Protein Concentration ( $A_{280}$ ):** The concentration of BSA is 20 mg/ml +/- 5% as determined by UV absorption at 280 nm. Protein concentration is determined by the Pace method using the extinction coefficient of 42,925 and molecular weight of 66,464 daltons for BSA (1).

**Non-Specific DNase Activity (16 hour):** A 50  $\mu$ l reaction in NEBuffer 4 containing 1  $\mu$ g of Lambda DNA (HindIII digest) and a minimum of 100  $\mu$ g of BSA incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.

**RNase Activity (2 Hour Digestion):** A 10  $\mu$ l reaction in NEBuffer 4 containing 40 ng of fluorescein labeled RNA transcript and a minimum of 20  $\mu$ g of BSA incubated for 2 hours at 37°C results in no detectable degradation of the RNA as determined by gel electrophoresis using fluorescence detection.

**RNase Activity (Extended Digestion):** A 10  $\mu$ l reaction in NEBuffer 4 containing 40 ng of fluorescein labeled RNA transcript and a minimum of 20  $\mu$ g of BSA is incubated at 37°C. After incubation for 16 hours, > 90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescence detection.

**Exonuclease Activity (Radioactivity Release):** A 50  $\mu$ l reaction in NEBuffer 4 containing 1  $\mu$ g of a mixture of single and double-stranded [ $^3$ H] *E. coli* DNA and a minimum of 100  $\mu$ g of BSA incubated for 4 hours at 37°C releases < 0.1% of the total radioactivity.

**Endonuclease Activity (Nicking):** A 50  $\mu$ l reaction in NEBuffer 4 containing 1  $\mu$ g of supercoiled  $\phi$ X174 RF I DNA and a minimum of 20  $\mu$ g of BSA incubated for 4 hours at 37°C results in < 20% conversion to the nicked form as determined by agarose gel electrophoresis.

**RNase Activity (Extended Digestion):** A 10  $\mu$ l reaction in NEBuffer 4 containing 40 ng of fluorescein labeled RNA transcript and a minimum of 20  $\mu$ g of BSA is incubated at 37°C. After incubation for 16 hours, > 90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescence detection.

**Exonuclease Activity (Radioactivity Release):** A 50  $\mu$ l reaction in NEBuffer 4 containing 1  $\mu$ g of a mixture of single and double-stranded [ $^3$ H] *E. coli* DNA and a minimum of 100  $\mu$ g of BSA incubated for 4 hours at 37°C releases < 0.1% of the total radioactivity.

**Endonuclease Activity (Nicking):** A 50  $\mu$ l reaction in NEBuffer 4 containing 1  $\mu$ g of supercoiled  $\phi$ X174 RF I DNA and a minimum of 20  $\mu$ g of BSA incubated for 4 hours at 37°C results in < 20% conversion to the nicked form as determined by agarose gel electrophoresis.

**Single Stranded DNase Activity (FAM Labeled Oligo):** A 50  $\mu$ l reaction in NEBuffer 4 containing a 20 nM solution of a fluorescent internal labeled oligonucleotide and a minimum of 100  $\mu$ g of BSA incubated for 16 hours at 37°C yields < 5% degradation as determined by capillary electrophoresis.

**Phosphatase Activity (FAM Labeled Oligo):** A 50  $\mu$ l reaction in NEBuffer 4 containing a 20 nM solution of a fluorescent internal labeled oligonucleotide with a 5' phosphate and a minimum of 100  $\mu$ g of BSA incubated for 16 hours at 37°C yields < 5% degradation as determined by capillary electrophoresis.

**qPCR DNA Contamination (*E. coli* Genomic):** A minimum of 10  $\mu$ g of BSA is screened for the presence of *E. coli* genomic DNA using SYBR Green qPCR with primers specific for the *E. coli* 16S rRNA locus. Results are quantified using a standard curve generated from purified *E. coli* genomic DNA. The measured level of *E. coli* genomic DNA contamination is less than 1 *E. coli* genome.

(see other side)

CERTIFICATE OF ANALYSIS

BSA,  
Molecular Biology  
Grade



1-800-632-7799  
info@neb.com  
www.neb.com



B9000S 006150618061

**B9000S**

**12 mg BSA, Molecular Biology Grade Exp: 6/18**  
**20 mg/ml Lot: 0061506 Store at -20°C**

**Description:** Bovine Serum Albumin (BSA) is supplied with some products to prevent adhesion of the enzyme to reaction tubes and pipette surfaces. BSA also stabilizes some proteins during incubation.

Supplied in: 20 mM Tris-HCl, 100 mM KCl, 0.1 mM EDTA and 50% glycerol (pH 8.0 @ 25°C).

**Single Stranded DNase Activity (FAM Labeled Oligo):** A 50  $\mu$ l reaction in NEBuffer 4 containing a 20 nM solution of a fluorescent internal labeled oligonucleotide and a minimum of 100  $\mu$ g of BSA incubated for 16 hours at 37°C yields < 5% degradation as determined by capillary electrophoresis.

**Phosphatase Activity (FAM Labeled Oligo):** A 50  $\mu$ l reaction in NEBuffer 4 containing a 20 nM solution of a fluorescent internal labeled oligonucleotide with a 5' phosphate and a minimum of 100  $\mu$ g of BSA incubated for 16 hours at 37°C yields < 5% degradation as determined by capillary electrophoresis.

**qPCR DNA Contamination (*E. coli* Genomic):** A minimum of 10  $\mu$ g of BSA is screened for the presence of *E. coli* genomic DNA using SYBR Green qPCR with primers specific for the *E. coli* 16S rRNA locus. Results are quantified using a standard curve generated from purified *E. coli* genomic DNA. The measured level of *E. coli* genomic DNA contamination is less than 1 *E. coli* genome.

(see other side)

CERTIFICATE OF ANALYSIS

**Reference:**

1. Pace, C.N. et al. (1995) *Protein Sci.*, 4, 2411–2423.



NEW ENGLAND BIOLABS® is a registered trademark of New England Biolabs, Inc.

This product is intended for research purposes only. This product is not intended to be used for therapeutic or diagnostic purposes in humans or animals.

**Reference:**

1. Pace, C.N. et al. (1995) *Protein Sci.*, 4, 2411–2423.



NEW ENGLAND BIOLABS® is a registered trademark of New England Biolabs, Inc.

This product is intended for research purposes only. This product is not intended to be used for therapeutic or diagnostic purposes in humans or animals.