

**$\beta$ -Nicotinamide  
adenine dinucleotide  
(NAD<sup>+</sup>)**



1-800-632-7799  
info@neb.com  
www.neb.com



B9007S 010141216121

**B9007S**

**0.2 ml 50 mM solution Lot: 0101412**

**Store at -20°C Exp: 12/16**

**Description:**  $\beta$ -Nicotinamide adenine dinucleotide (NAD<sup>+</sup>) is included with the PreCR™ Repair Mix available from New England Biolabs. NAD<sup>+</sup> is supplied as an aqueous 50 mM solution.

**Use and Storage:** For use in the PreCR Repair Mix, NAD<sup>+</sup> should be diluted 1:100 to a final concentration of 500  $\mu$ M. NAD<sup>+</sup> solution should be stored at -20°C, and the user should avoid repeated freeze/thaw cycles. For storage longer than 30 days, store at -80°C.

CERTIFICATE OF ANALYSIS

**$\beta$ -Nicotinamide  
adenine dinucleotide  
(NAD<sup>+</sup>)**



1-800-632-7799  
info@neb.com  
www.neb.com



B9007S 010141216121

**B9007S**

**0.2 ml 50 mM solution Lot: 0101412**

**Store at -20°C Exp: 12/16**

**Description:**  $\beta$ -Nicotinamide adenine dinucleotide (NAD<sup>+</sup>) is included with the PreCR™ Repair Mix available from New England Biolabs. NAD<sup>+</sup> is supplied as an aqueous 50 mM solution.

**Use and Storage:** For use in the PreCR Repair Mix, NAD<sup>+</sup> should be diluted 1:100 to a final concentration of 500  $\mu$ M. NAD<sup>+</sup> solution should be stored at -20°C, and the user should avoid repeated freeze/thaw cycles. For storage longer than 30 days, store at -80°C.

CERTIFICATE OF ANALYSIS