

HhaI Methyltransferase



1-800-632-7799
info@neb.com
www.neb.com




M0217S 006120513051

M0217S



1,000 units 25,000 U/ml Lot: 0061205
RECOMBINANT Store at -20°C Exp: 5/13

Methylation Site:



Description: HhaI Methyltransferase modifies the internal cytosine residue (C⁵) in the sequence above.

HhaI Methyltransferase



1-800-632-7799
info@neb.com
www.neb.com




M0217S 006120513051

M0217S



1,000 units 25,000 U/ml Lot: 0061205
RECOMBINANT Store at -20°C Exp: 5/13

Methylation Site:



Description: HhaI Methyltransferase modifies the internal cytosine residue (C⁵) in the sequence above.

Source: An *E. coli* strain that carries the cloned HhaI modification gene from *Haemophilus haemolyticus* (ATCC 10014)

Supplied in: 150 mM NaCl, 50 mM Tris-HCl (pH 7.5), 10 mM EDTA, 5 mM 2-mercapto-ethanol, 200 µg/ml BSA and 50% glycerol.

Reagents Supplied with Enzyme:

10X HhaI Methyltransferase Reaction Buffer,
400X S-adenosylmethionine (32 mM).

Reaction Conditions:

1X HhaI Methyltransferase Reaction Buffer, 80 µM S-adenosylmethionine.
Incubate at 37°C.

1X HhaI Methyltransferase Reaction Buffer:

50 mM Tris-HCl
10 mM EDTA
5 mM 2-mercaptoethanol
pH 7.5 @ 25°C

Protection Assay Conditions:

HhaI Methyltransferase is incubated with 1 µg of λ DNA in 10 µl 1X HhaI Methyltransferase Reaction Buffer, supplemented with 80 µM S-adenosylmethionine, for one hour at 37°C followed by 15 minutes at 65°C.

Source: An *E. coli* strain that carries the cloned HhaI modification gene from *Haemophilus haemolyticus* (ATCC 10014)

Supplied in: 150 mM NaCl, 50 mM Tris-HCl (pH 7.5), 10 mM EDTA, 5 mM 2-mercapto-ethanol, 200 µg/ml BSA and 50% glycerol.

Reagents Supplied with Enzyme:

10X HhaI Methyltransferase Reaction Buffer,
400X S-adenosylmethionine (32 mM).

Reaction Conditions:

1X HhaI Methyltransferase Reaction Buffer, 80 µM S-adenosylmethionine.
Incubate at 37°C.

1X HhaI Methyltransferase Reaction Buffer:

50 mM Tris-HCl
10 mM EDTA
5 mM 2-mercaptoethanol
pH 7.5 @ 25°C

Protection Assay Conditions:

HhaI Methyltransferase is incubated with 1 µg of λ DNA in 10 µl 1X HhaI Methyltransferase Reaction Buffer, supplemented with 80 µM S-adenosylmethionine, for one hour at 37°C followed by 15 minutes at 65°C.

The extent of protection by HhaI Methyltransferase is determined by the addition of 40 µl NEBuffer 4 supplemented with 100 µg/ml BSA, 10 mM MgCl₂ and 10 units of Hha I restriction endonuclease. Incubation at 37°C for 30 minutes is followed by analysis on an agarose gel.

Unit Definition:

One unit is defined as the amount of enzyme required to protect 1 µg λ DNA in 1 hour at 37°C in a total reaction volume of 10 µl against cleavage by Hha I restriction endonuclease.

Quality Control Assays

16-Hour Incubation: Incubation of 125 units of HhaI Methyltransferase with 1 µg of HindIII-digested DNA in 50 µl 1X NEBuffer 2 for 16 hours at 37°C resulted in no detectable endonuclease contamination.

Exonuclease Activity: Incubation of 250 units of HhaI Methyltransferase with 1 µg sonicated ³H DNA (10⁵ cpm/µg) for 4 hours at 37°C in 50 µl NEBuffer 2 [50 mM NaCl, 10 mM Tris-HCl (pH 7.9 @ 25°C), 10 mM MgCl₂, 1 mM DTT] released 0.39% of the total radioactivity.

The extent of protection by HhaI Methyltransferase is determined by the addition of 40 µl NEBuffer 4 supplemented with 100 µg/ml BSA, 10 mM MgCl₂ and 10 units of Hha I restriction endonuclease. Incubation at 37°C for 30 minutes is followed by analysis on an agarose gel.

Unit Definition:

One unit is defined as the amount of enzyme required to protect 1 µg λ DNA in 1 hour at 37°C in a total reaction volume of 10 µl against cleavage by Hha I restriction endonuclease.

Quality Control Assays

16-Hour Incubation: Incubation of 125 units of HhaI Methyltransferase with 1 µg of HindIII-digested DNA in 50 µl 1X NEBuffer 2 for 16 hours at 37°C resulted in no detectable endonuclease contamination.

Exonuclease Activity: Incubation of 250 units of HhaI Methyltransferase with 1 µg sonicated ³H DNA (10⁵ cpm/µg) for 4 hours at 37°C in 50 µl NEBuffer 2 [50 mM NaCl, 10 mM Tris-HCl (pH 7.9 @ 25°C), 10 mM MgCl₂, 1 mM DTT] released 0.39% of the total radioactivity.

Storage of SAM: S-adenosylmethionine or SAM is stored at -20°C as a 32 mM solution dissolved in 0.005 M sulfuric acid and 10% ethanol. Under these conditions SAM is stable for up to 6 months. SAM is unstable at (pH 7.5), 37°C, (1) and should be replenished in reactions incubated longer than 4 hours.

Methylation can be optimized by using fresh SAM.

Reference:

- Hoffman, J. L. (1986) *Biochemistry* 25, 4444-4449.

Companion Product:

S-adenosylmethionine (SAM)
#B9003S 0.5 ml

CERTIFICATE OF ANALYSIS

Storage of SAM: S-adenosylmethionine or SAM is stored at -20°C as a 32 mM solution dissolved in 0.005 M sulfuric acid and 10% ethanol. Under these conditions SAM is stable for up to 6 months. SAM is unstable at (pH 7.5), 37°C, (1) and should be replenished in reactions incubated longer than 4 hours.

Methylation can be optimized by using fresh SAM.

Reference:

- Hoffman, J. L. (1986) *Biochemistry* 25, 4444-4449.

Companion Product:

S-adenosylmethionine (SAM)
#B9003S 0.5 ml

CERTIFICATE OF ANALYSIS