

HaeIII Methyltransferase



1-800-632-7799
info@neb.com
www.neb.com



M0224S 006151016101


M0224S



500 units 10,000 U/ml Lot: 0061510

RECOMBINANT Store at -20°C Exp: 10/16

Methylation Site:



Description: HaeIII Methyltransferase modifies the internal cytosine residue (C⁵) in the sequence above.

HaeIII Methyltransferase



1-800-632-7799
info@neb.com
www.neb.com



M0224S 006151016101


M0224S



500 units 10,000 U/ml Lot: 0061510

RECOMBINANT Store at -20°C Exp: 10/16

Methylation Site:



Description: HaeIII Methyltransferase modifies the internal cytosine residue (C⁵) in the sequence above.

Source: An *E. coli* strain that carries the cloned HaeIII modification gene from *Haemophilus aegyptius* (ATCC 11116)

Supplied in: 50 mM KCl, 50 mM Tris-HCl (pH 7.5), 10 mM EDTA, 1 mM dithiothreitol, 200 µg/ml BSA and 50% glycerol.

Reagents Supplied with Enzyme:

10X HaeIII Methyltransferase Reaction Buffer, 400X S-adenosylmethionine (32 mM).

Reaction Conditions:

1X HaeIII Methyltransferase Reaction Buffer, supplemented with 80 µM S-adenosylmethionine (supplied). Incubate at 37°C.

1X HaeIII Methyltransferase Reaction Buffer:

50 mM NaCl
50 mM Tris-HCl
10 mM dithiothreitol
pH 8.5 @ 25°C

Protection Assay Conditions:

HaeIII Methyltransferase is incubated with 1 µg of λ DNA in 10 µl of 1X HaeIII Methyltransferase Reaction Buffer, supplemented

Source: An *E. coli* strain that carries the cloned HaeIII modification gene from *Haemophilus aegyptius* (ATCC 11116)

Supplied in: 50 mM KCl, 50 mM Tris-HCl (pH 7.5), 10 mM EDTA, 1 mM dithiothreitol, 200 µg/ml BSA and 50% glycerol.

Reagents Supplied with Enzyme:

10X HaeIII Methyltransferase Reaction Buffer, 400X S-adenosylmethionine (32 mM).

Reaction Conditions:

1X HaeIII Methyltransferase Reaction Buffer, supplemented with 80 µM S-adenosylmethionine (supplied). Incubate at 37°C.

1X HaeIII Methyltransferase Reaction Buffer:

50 mM NaCl
50 mM Tris-HCl
10 mM dithiothreitol
pH 8.5 @ 25°C

Protection Assay Conditions:

HaeIII Methyltransferase is incubated with 1 µg of λ DNA in 10 µl of 1X HaeIII Methyltransferase Reaction Buffer, supplemented

with 80 µM S-adenosylmethionine, for one hour at 37°C followed by 15 minutes at 65°C. The extent of protection is determined by the addition of 40 µl NEBuffer 2 and 10 units of HaeIII restriction endonuclease. Incubation for 1 hour at 37°C is followed by analysis on an agarose gel.

Unit Definition:

One unit is defined as the amount of enzyme required to protect 1 µg λ DNA in 1 hour at 37°C in a total reaction volume of 10 µl against cleavage by HaeIII restriction endonuclease.

Quality Control Assays

16-Hour Incubation: Incubation of 100 units with 1 µg of HindIII-digested λ DNA in 50 µl 1X NEBuffer 2 for 16 hours at 37°C resulted in no detectable endonuclease contamination.

Exonuclease Activity: Incubation of 100 units of HaeIII Methyltransferase with 1 µg sonicated ³H-DNA (10⁵ cpm/µg) for 4 hours at 37°C in 50 µl NEBuffer 2 [50 mM NaCl, 10 mM Tris-HCl (pH 7.9 @ 25°C), 10 mM MgCl₂, 1 mM dithiothreitol] released < 0.05% of the total radioactivity.

with 80 µM S-adenosylmethionine, for one hour at 37°C followed by 15 minutes at 65°C. The extent of protection is determined by the addition of 40 µl NEBuffer 2 and 10 units of HaeIII restriction endonuclease. Incubation for 1 hour at 37°C is followed by analysis on an agarose gel.

Unit Definition:

One unit is defined as the amount of enzyme required to protect 1 µg λ DNA in 1 hour at 37°C in a total reaction volume of 10 µl against cleavage by HaeIII restriction endonuclease.

Quality Control Assays

16-Hour Incubation: Incubation of 100 units with 1 µg of HindIII-digested λ DNA in 50 µl 1X NEBuffer 2 for 16 hours at 37°C resulted in no detectable endonuclease contamination.

Exonuclease Activity: Incubation of 100 units of HaeIII Methyltransferase with 1 µg sonicated ³H-DNA (10⁵ cpm/µg) for 4 hours at 37°C in 50 µl NEBuffer 2 [50 mM NaCl, 10 mM Tris-HCl (pH 7.9 @ 25°C), 10 mM MgCl₂, 1 mM dithiothreitol] released < 0.05% of the total radioactivity.

Endonuclease Contamination: Incubation of 100 units with 1 µg φX174 RF I DNA (4 hours, 37°C in 50 µl of NEBuffer 2) gave < 5% conversion to RF II.

Storage of SAM: S-adenosylmethionine (SAM) is stored at -20°C as a 32 mM solution dissolved in 0.005 M sulfuric acid and 10% ethanol. Under these conditions SAM is stable for up to 6 months. SAM is unstable at (pH 7.5), 37°C (1) and should be replenished in reactions incubated longer than 4 hours.

Methylation can be optimized by using fresh SAM.

Note: HaeIII Methyltransferase protects DNA against cleavage by NotI.

Reference:

- Hoffman, J.L. (1986) *Biochemistry* 25, 4444-4449.

Companion Product:

S-adenosylmethionine (SAM)
#B9003S 0.5 ml

CERTIFICATE OF ANALYSIS

Endonuclease Contamination: Incubation of 100 units with 1 µg φX174 RF I DNA (4 hours, 37°C in 50 µl of NEBuffer 2) gave < 5% conversion to RF II.

Storage of SAM: S-adenosylmethionine (SAM) is stored at -20°C as a 32 mM solution dissolved in 0.005 M sulfuric acid and 10% ethanol. Under these conditions SAM is stable for up to 6 months. SAM is unstable at (pH 7.5), 37°C (1) and should be replenished in reactions incubated longer than 4 hours.

Methylation can be optimized by using fresh SAM.

Note: HaeIII Methyltransferase protects DNA against cleavage by NotI.

Reference:

- Hoffman, J.L. (1986) *Biochemistry* 25, 4444-4449.

Companion Product:

S-adenosylmethionine (SAM)
#B9003S 0.5 ml

CERTIFICATE OF ANALYSIS



NEW ENGLAND BIOLABS® is a registered trademark of New England Biolabs, Inc.

This product is intended for research purposes only. This product is not intended to be used for therapeutic or diagnostic purposes in humans or animals.



NEW ENGLAND BIOLABS® is a registered trademark of New England Biolabs, Inc.

This product is intended for research purposes only. This product is not intended to be used for therapeutic or diagnostic purposes in humans or animals.