

SET7 Methyltransferase



M0233S 004160117011

M0233S



100 units **2,000 U/ml** **Lot: 0041601**
RECOMBINANT **Store at -20°C** **Exp: 11/17**

Description: SET7 Methyltransferase methylates lysine 4 (Lys 4) of histone H3 (1). Methylation occurs at the ε amino group of lysine residues (2,3). Di- and tri- methylation of histone H3 Lys 4 is a hallmark of transcriptionally active chromatin and is globally distributed (4,5).

SET7 Methyltransferase



M0233S 004160117011

M0233S



100 units **2,000 U/ml** **Lot: 0041601**
RECOMBINANT **Store at -20°C** **Exp: 11/17**

Description: SET7 Methyltransferase methylates lysine 4 (Lys 4) of histone H3 (1). Methylation occurs at the ε amino group of lysine residues (2,3). Di- and tri- methylation of histone H3 Lys 4 is a hallmark of transcriptionally active chromatin and is globally distributed (4,5).

Source: SET7 enzyme is expressed from human SET7 cDNA using an *E. coli* expression system using an MBP fusion tag.

Supplied in: 50 mM Tris-HCl (pH 8.0 @ 25°C), 5 mM MgCl₂, 100 mM NaCl, 4 mM dithiothreitol and 50% glycerol.

Reagents Supplied with Enzyme:
10X HMT Reaction Buffer
32 mM S-adenosylmethionine (SAM)

Reaction Conditions: 1X HMT Reaction Buffer supplemented with 160 μM S-adenosylmethionine. Incubate at 37°C.

1X HMT Reaction Buffer:
50 mM Tris-HCl
5 mM MgCl₂
4 mM dithiothreitol
(pH 9.0 @25°C)

Source: SET7 enzyme is expressed from human SET7 cDNA using an *E. coli* expression system using an MBP fusion tag.

Supplied in: 50 mM Tris-HCl (pH 8.0 @ 25°C), 5 mM MgCl₂, 100 mM NaCl, 4 mM dithiothreitol and 50% glycerol.

Reagents Supplied with Enzyme:
10X HMT Reaction Buffer
32 mM S-adenosylmethionine (SAM)

Reaction Conditions: 1X HMT Reaction Buffer supplemented with 160 μM S-adenosylmethionine. Incubate at 37°C.

1X HMT Reaction Buffer:
50 mM Tris-HCl
5 mM MgCl₂
4 mM dithiothreitol
(pH 9.0 @25°C)

Unit Definition: One unit is defined as the amount of enzyme required to catalyze the transfer of 50 pmol of methyl group to substrate histone H3 in a total reaction volume of 25 μl in 10 minutes at 37°C.

Quality Assurance: Purified free of contaminating proteases.

Storage Note: S-adenosylmethionine (SAM) is stored at -20°C as a 32 mM solution dissolved in 0.005 M sulfuric acid and 10% ethanol (pH 7.5). Under these conditions, SAM is stable for up to 6 months. SAM is unstable at 37°C and should be replenished in reactions incubated longer than 4 hours. Methylation can be optimized by using fresh SAM.

Heat Inactivation: 65°C for 20 minutes.

Unit Definition: One unit is defined as the amount of enzyme required to catalyze the transfer of 50 pmol of methyl group to substrate histone H3 in a total reaction volume of 25 μl in 10 minutes at 37°C.

Quality Assurance: Purified free of contaminating proteases.

Storage Note: S-adenosylmethionine (SAM) is stored at -20°C as a 32 mM solution dissolved in 0.005 M sulfuric acid and 10% ethanol (pH 7.5). Under these conditions, SAM is stable for up to 6 months. SAM is unstable at 37°C and should be replenished in reactions incubated longer than 4 hours. Methylation can be optimized by using fresh SAM.

Heat Inactivation: 65°C for 20 minutes.

References:

1. Wang, H. et al. (2001) *Mol Cell*, 6, 1207–1217.
2. Xiao, B. et al. (2003) *Nature*, 421, 652–656.
3. Wilson, J.R. et al. (2002) *Cell*, 111, 105–115.
4. Schneider, R. et al. (2004) *Nat. Cell Biol.* 6, 73–77.
5. Santos-Rosa, H. et al. (2002) *Nature*, 419, 407–411.



NEW ENGLAND BIOLABS® is a registered trademark of New England Biolabs, Inc.

This product is intended for research purposes only. This product is not intended to be used for therapeutic or diagnostic purposes in humans or animals.

CERTIFICATE OF ANALYSIS

References:

1. Wang, H. et al. (2001) *Mol Cell*, 6, 1207–1217.
2. Xiao, B. et al. (2003) *Nature*, 421, 652–656.
3. Wilson, J.R. et al. (2002) *Cell*, 111, 105–115.
4. Schneider, R. et al. (2004) *Nat. Cell Biol.* 6, 73–77.
5. Santos-Rosa, H. et al. (2002) *Nature*, 419, 407–411.



NEW ENGLAND BIOLABS® is a registered trademark of New England Biolabs, Inc.

This product is intended for research purposes only. This product is not intended to be used for therapeutic or diagnostic purposes in humans or animals.

CERTIFICATE OF ANALYSIS