



M0240S



500 units 8,000 U/ml Lot: 0081605 RECOMBINANT Store at -20°C Exp: 5/18

Description: Fpg (formamidopyrimidine [fapy]-DNA glycosylase) (also known as 8-oxoguanine DNA glycosylase) acts both as a N-glycosylase and an AP-lyase. The N-glycosylase activity releases damaged purines from double-stranded DNA, generating an apurinic (AP site). The AP-lyase activity cleaves both 3' and 5' to the AP site thereby removing the AP site and leaving a 1 base gap with a 5' and 3' phosphate.

Some of the damaged bases recognized and



M0240S



500 units 8,000 U/ml Lot: 0081605 RECOMBINANT Store at -20°C Exp: 5/18

Description: Fpg (formamidopyrimidine [fapy]-DNA glycosylase) (also known as 8-oxoguanine DNA glycosylase) acts both as a N-glycosylase and an AP-lyase. The N-glycosylase activity releases damaged purines from double-stranded DNA, generating an apurinic (AP site). The AP-lyase activity cleaves both 3' and 5' to the AP site thereby removing the AP site and leaving a 1 base gap with a 5' and 3' phosphate.

Some of the damaged bases recognized and

removed by Fpg include 7, 8-dihydro-8-oxoguanine (8-oxoguanine), 8-oxoadenine, fapy-guanine, methy-fapy-guanine, fapy-adenine, aflatoxin B1-fapy-guanine, 5-hydroxy-cytosine and 5-hydroxy-uracil (1,2).

Source: An E. coli strain that carries the cloned fpg gene (3)

Applications:

- Single cell gel electrophoresis (Comet assay) (4,5,6)
Alkaline elution (7)
Alkaline unwinding (8)
Modified nick translation (9)

Supplied in: 20 mM Tris-HCl (pH 8.0), 0.5 mM EDTA, 50 mM NaCl, 200 µg/ml BSA and 50% glycerol.

Reagents Supplied with Enzyme: 10X NEBuffer 1, 100X BSA.

Reaction Conditions: 1X NEBuffer 1,

supplemented with 100 µg/ml BSA. Incubate at 37°C.

1X NEBuffer 1:

10 mM Bis Tris Propane-HCl
10 mM MgCl2
1 mM DTT
pH 7.0 @ 25°C

When using a buffer other than the optimal (supplied) NEBuffer, it may be necessary to add more enzyme to achieve complete digestion.

Unit Definition: One unit is defined as the amount of enzyme required to cleave 1 pmol of a 34-mer oligonucleotide duplex containing a single 8-oxoguanine base paired with a cytosine in a total reaction volume of 10 µl in 1 hour at 37°C.

Unit Assay Conditions: 1X NEBuffer 1 containing 10 pmol of fluorescently labeled oligonucleotide duplex, supplemented with 100 µg/ml BSA in a total reaction volume of 10 µl.

Recommended Dilution for the Comet Assay: 1:10^3 to 1:10^4 (4,5,6,10). A detailed protocol can be found at www.neb.com.

supplemented with 100 µg/ml BSA. Incubate at 37°C.

1X NEBuffer 1:

10 mM Bis Tris Propane-HCl
10 mM MgCl2
1 mM DTT
pH 7.0 @ 25°C

When using a buffer other than the optimal (supplied) NEBuffer, it may be necessary to add more enzyme to achieve complete digestion.

Unit Definition: One unit is defined as the amount of enzyme required to cleave 1 pmol of a 34-mer oligonucleotide duplex containing a single 8-oxoguanine base paired with a cytosine in a total reaction volume of 10 µl in 1 hour at 37°C.

Unit Assay Conditions: 1X NEBuffer 1 containing 10 pmol of fluorescently labeled oligonucleotide duplex, supplemented with 100 µg/ml BSA in a total reaction volume of 10 µl.

Recommended Dilution for the Comet Assay: 1:10^3 to 1:10^4 (4,5,6,10). A detailed protocol can be found at www.neb.com.

Quality Control Assays

16-Hour Incubation: A 50 µl reaction containing 1 µg of λ DNA (HindIII digest) and 40 units of Fpg incubated for 16 hours at 37°C resulted in DNA patterns free of detectable nuclease degradation as determined by agarose gel electrophoresis.

Exonuclease Activity: Incubation of a 50 µl reaction containing 40 units of Fpg with 1 µg of a mixture of single and double-stranded [3H] E. coli DNA (10^5 cpm/µg) for 4 hours at 37°C released < 1.0% of the total radioactivity.

Heat Inactivation: 160 units of enzyme were inactivated by incubation at 60°C for 10 minutes.

Physical Purity: Purified to > 95% homogeneity as determined by SDS-PAGE analysis using Coomassie Blue detection. BSA is added to the enzyme for stability.

Usage Note: Fpg will remove deoxyribose- 5' phosphate dR5'P at a nicked site (11).

(see other side)

CERTIFICATE OF ANALYSIS

Quality Control Assays

16-Hour Incubation: A 50 µl reaction containing 1 µg of λ DNA (HindIII digest) and 40 units of Fpg incubated for 16 hours at 37°C resulted in DNA patterns free of detectable nuclease degradation as determined by agarose gel electrophoresis.

Exonuclease Activity: Incubation of a 50 µl reaction containing 40 units of Fpg with 1 µg of a mixture of single and double-stranded [3H] E. coli DNA (10^5 cpm/µg) for 4 hours at 37°C released < 1.0% of the total radioactivity.

Heat Inactivation: 160 units of enzyme were inactivated by incubation at 60°C for 10 minutes.

Physical Purity: Purified to > 95% homogeneity as determined by SDS-PAGE analysis using Coomassie Blue detection. BSA is added to the enzyme for stability.

Usage Note: Fpg will remove deoxyribose- 5' phosphate dR5'P at a nicked site (11).

(see other side)

CERTIFICATE OF ANALYSIS

References:

1. Tchou, J., et al. (1994). Substrate specificity of Fpg protein. *J. Biol. Chem.* 269, 1518–1524.
2. Hatahet, Z., et al. (1994). New substrates for old enzymes. *J. Biol Chem.* 269, 18814–18820.
3. Boiteux, S., O'Connor, T. and Laval, J. (1987). Formamidopyrimidine-DNA glycosylase of *Escherichia coli*: cloning and sequencing of the *fpg* structural gene and overproduction of the protein. *EMBO J.* 5, 3177–3183.
4. Singh, N., McCoy, M., Tice, R. and Schneider, L. (1988). A simple technique for quantitation of low levels of DNA damage in individual cells. *Experimental Cell Research* 175, 184–191.
5. Collins, A., Duthie, S. and Dobson, V. (1993). Direct enzymatic detection of endogenous oxidative base damage in human lymphocyte DNA. *Carcinogenesis* 14, 1733–1735.
6. Collins, A., Dusinska, M., Gedik, C. and Stetina, R. (1996). Oxidative damage to DNA: do we have a reliable biomarker? *Environmental Health Perspectives* 104, 465–469.
7. Pflaum, M., Will, O., Mahler, H-C. and Epe, B. (1998). DNA oxidation products determined with repair endonucleases in mammalian cells: types, basal levels and influence of cell proliferation. *Free Rad. Res.* 29, 585–594
8. Hartwig, A., Dally, H. and Schlepegrell, R. (1996). Sensitive analysis of oxidative DNA damage in mammalian cells: use of the bacterial Fpg protein in combination with alkaline unwinding. *Toxicology Letters* 88, 85–90.
9. Czene, S. and Harms-Ringdahl, M. (1995). Detection of single strand breaks and formamidopyrimidine-DNA glycosylase-sensitive sites in DNA of cultured human fibroblasts. *Mutation Research* 336, 235–242.
10. Guthrie, E., New England Biolabs, Inc., unpublished observations.
11. Marks, K. and Landry D., New England Biolabs, Inc., unpublished observations.

Page 2 (M0240)

References:

1. Tchou, J., et al. (1994). Substrate specificity of Fpg protein. *J. Biol. Chem.* 269, 1518–1524.
2. Hatahet, Z., et al. (1994). New substrates for old enzymes. *J. Biol Chem.* 269, 18814–18820.
3. Boiteux, S., O'Connor, T. and Laval, J. (1987). Formamidopyrimidine-DNA glycosylase of *Escherichia coli*: cloning and sequencing of the *fpg* structural gene and overproduction of the protein. *EMBO J.* 5, 3177–3183.
4. Singh, N., McCoy, M., Tice, R. and Schneider, L. (1988). A simple technique for quantitation of low levels of DNA damage in individual cells. *Experimental Cell Research* 175, 184–191.
5. Collins, A., Duthie, S. and Dobson, V. (1993). Direct enzymatic detection of endogenous oxidative base damage in human lymphocyte DNA. *Carcinogenesis* 14, 1733–1735.
6. Collins, A., Dusinska, M., Gedik, C. and Stetina, R. (1996). Oxidative damage to DNA: do we have a reliable biomarker? *Environmental Health Perspectives* 104, 465–469.
7. Pflaum, M., Will, O., Mahler, H-C. and Epe, B. (1998). DNA oxidation products determined with repair endonucleases in mammalian cells: types, basal levels and influence of cell proliferation. *Free Rad. Res.* 29, 585–594
8. Hartwig, A., Dally, H. and Schlepegrell, R. (1996). Sensitive analysis of oxidative DNA damage in mammalian cells: use of the bacterial Fpg protein in combination with alkaline unwinding. *Toxicology Letters* 88, 85–90.
9. Czene, S. and Harms-Ringdahl, M. (1995). Detection of single strand breaks and formamidopyrimidine-DNA glycosylase-sensitive sites in DNA of cultured human fibroblasts. *Mutation Research* 336, 235–242.
10. Guthrie, E., New England Biolabs, Inc., unpublished observations.
11. Marks, K. and Landry D., New England Biolabs, Inc., unpublished observations.

Page 2 (M0240)



NEW ENGLAND BIOLABS® is a registered trademark of New England Biolabs, Inc.

This product is intended for research purposes only. This product is not intended to be used for therapeutic or diagnostic purposes in humans or animals.



NEW ENGLAND BIOLABS® is a registered trademark of New England Biolabs, Inc.

This product is intended for research purposes only. This product is not intended to be used for therapeutic or diagnostic purposes in humans or animals.