

Mung Bean Nuclease



1-800-632-7799
info@neb.com
www.neb.com



M0250S 025131015101

M0250S

1,500 units **Lot: 0251310** **Exp: 10/15**
10,000 U/ml **Store at -20°C**

Description: A single-strand specific DNA and RNA endonuclease which will degrade single-stranded extensions from the ends of DNA and RNA molecules, leaving blunt, ligatable ends.

Source: Mung bean sprouts

Molecular Weight: 39 kDa

Supplied in: 10 mM sodium acetate (pH 5.0)
0.1 mM zinc acetate, 1 mM cysteine, 0.001%
Triton X-100 and 50% glycerol.

Applications:

- Removal of 3' and 5' extensions from DNA or RNA termini
- Transcriptional mapping
- Cleavage of hairpin loops
- Excision of gene coding sequences from genomic DNA
- Generation of new restriction sites

Note: It is no longer necessary to supplement Mung Bean Nuclease reactions with Zn²⁺. The zinc acetate in the storage buffer fulfills the Zn²⁺ requirement of the enzyme even after dilution in a reaction.

Reagents Supplied with Enzyme:

10X Mung Bean Nuclease Reaction Buffer

Reaction Conditions: Substrate DNA at a concentration of 0.1 µg/µl in 1X Mung Bean Nuclease Reaction Buffer.

Incubate at 30°C.

Applications:

- Removal of 3' and 5' extensions from DNA or RNA termini
- Transcriptional mapping
- Cleavage of hairpin loops
- Excision of gene coding sequences from genomic DNA
- Generation of new restriction sites

Note: It is no longer necessary to supplement Mung Bean Nuclease reactions with Zn²⁺. The zinc acetate in the storage buffer fulfills the Zn²⁺ requirement of the enzyme even after dilution in a reaction.

Reagents Supplied with Enzyme:

10X Mung Bean Nuclease Reaction Buffer

Reaction Conditions: Substrate DNA at a concentration of 0.1 µg/µl in 1X Mung Bean Nuclease Reaction Buffer.

Incubate at 30°C.

1X Mung Bean Nuclease Reaction Buffer:

50 mM sodium acetate
30 mM NaCl
1 mM ZnSO₄
pH 5.0 @ 25°C

Also active in NEBuffers 1, 2 & 4.

Unit Definition: One unit is defined as the amount of enzyme required to produce 1 µg of acid-soluble total nucleotide in 1 minute at 37°C.

Unit Assay Conditions: 1X Mung Bean Nuclease Reaction Buffer and 0.5 mg/ml denatured calf thymus DNA as an enzyme substrate.

Removal of Single-Stranded Extensions:

1. Suspend DNA (0.1 µg/µl) in 1X Mung Bean Nuclease Reaction Buffer or 1X NEBuffers 1, 2, or 4.
2. Add 1.0 unit of Mung Bean Nuclease per µg DNA.
3. Incubate at 30°C for 30 minutes.
4. Inactivate the enzyme by phenol/chloroform extraction or by addition of SDS to 0.01%.
5. Recover the DNA by ethanol precipitation.

1X Mung Bean Nuclease Reaction Buffer:

50 mM sodium acetate
30 mM NaCl
1 mM ZnSO₄
pH 5.0 @ 25°C

Also active in NEBuffers 1, 2 & 4.

Unit Definition: One unit is defined as the amount of enzyme required to produce 1 µg of acid-soluble total nucleotide in 1 minute at 37°C.

Unit Assay Conditions: 1X Mung Bean Nuclease Reaction Buffer and 0.5 mg/ml denatured calf thymus DNA as an enzyme substrate.

Removal of Single-Stranded Extensions:

1. Suspend DNA (0.1 µg/µl) in 1X Mung Bean Nuclease Reaction Buffer or 1X NEBuffers 1, 2, or 4.
2. Add 1.0 unit of Mung Bean Nuclease per µg DNA.
3. Incubate at 30°C for 30 minutes.
4. Inactivate the enzyme by phenol/chloroform extraction or by addition of SDS to 0.01%.
5. Recover the DNA by ethanol precipitation.

Quality Assurance: Purified free of double-strand exonuclease contamination.

Quality Control Assays

16 µg of Hae III digested φX174 DNA was incubated with 10 units of Mung Bean Nuclease in a 400 µl volume of 1X NEBuffer 2 for 30 minutes at 30°C. The DNA was then precipitated, ligated with T4 DNA Ligase and recut. 90% of the DNA fragments treated with Mung Bean Nuclease were ligated and of those 95% were recut with Hae III.

References:

1. Kowalski, D. et al. (1976) *Biochemistry* 15, 4457-4463.
2. McCutchan, T.F. et al. (1984) *Science* 225, 626-628.

CERTIFICATE OF ANALYSIS

Mung Bean Nuclease



1-800-632-7799
info@neb.com
www.neb.com



M0250S 025131015101

M0250S

1,500 units **Lot: 0251310** **Exp: 10/15**
10,000 U/ml **Store at -20°C**

Description: A single-strand specific DNA and RNA endonuclease which will degrade single-stranded extensions from the ends of DNA and RNA molecules, leaving blunt, ligatable ends.

Source: Mung bean sprouts

Molecular Weight: 39 kDa

Supplied in: 10 mM sodium acetate (pH 5.0)
0.1 mM zinc acetate, 1 mM cysteine, 0.001%
Triton X-100 and 50% glycerol.

Quality Assurance: Purified free of double-strand exonuclease contamination.

Quality Control Assays

16 µg of Hae III digested φX174 DNA was incubated with 10 units of Mung Bean Nuclease in a 400 µl volume of 1X NEBuffer 2 for 30 minutes at 30°C. The DNA was then precipitated, ligated with T4 DNA Ligase and recut. 90% of the DNA fragments treated with Mung Bean Nuclease were ligated and of those 95% were recut with Hae III.

References:

1. Kowalski, D. et al. (1976) *Biochemistry* 15, 4457-4463.
2. McCutchan, T.F. et al. (1984) *Science* 225, 626-628.

CERTIFICATE OF ANALYSIS