

Mung Bean Nuclease



1-800-632-7799
info@neb.com
www.neb.com



M0250S 025160818081

M0250S

1,500 units **Lot: 0251608** **Exp: 8/18**
10,000 U/ml **Store at -20°C**

Description: A single-strand specific DNA and RNA endonuclease which will degrade single-stranded extensions from the ends of DNA and RNA molecules, leaving blunt, ligatable ends.

Source: Mung bean sprouts

Molecular Weight: 39 kDa

Supplied in: 10 mM sodium acetate (pH 5.0)
0.1 mM zinc acetate, 1 mM cysteine, 0.001%
Triton X-100 and 50% glycerol.

Applications:

- Removal of 3' and 5' extensions from DNA or RNA termini
- Transcriptional mapping
- Cleavage of hairpin loops
- Excision of gene coding sequences from genomic DNA
- Generation of new restriction sites

Note: It is no longer necessary to supplement Mung Bean Nuclease reactions with Zn²⁺. The zinc acetate in the storage buffer fulfills the Zn²⁺ requirement of the enzyme even after dilution in a reaction.

Reagents Supplied with Enzyme:
10X Mung Bean Nuclease Reaction Buffer

Reaction Conditions: Substrate DNA at a concentration of 0.1 µg/µl in 1X Mung Bean Nuclease Reaction Buffer. **Incubate at 30°C.**

Applications:

- Removal of 3' and 5' extensions from DNA or RNA termini
- Transcriptional mapping
- Cleavage of hairpin loops
- Excision of gene coding sequences from genomic DNA
- Generation of new restriction sites

Note: It is no longer necessary to supplement Mung Bean Nuclease reactions with Zn²⁺. The zinc acetate in the storage buffer fulfills the Zn²⁺ requirement of the enzyme even after dilution in a reaction.

Reagents Supplied with Enzyme:
10X Mung Bean Nuclease Reaction Buffer

Reaction Conditions: Substrate DNA at a concentration of 0.1 µg/µl in 1X Mung Bean Nuclease Reaction Buffer. **Incubate at 30°C.**

1X Mung Bean Nuclease Reaction Buffer:

50 mM sodium acetate
30 mM NaCl
1 mM ZnSO₄
pH 5.0 @ 25°C
Also active in NEBuffers 1.1, 2.1 or CutSmart®.

Unit Definition: One unit is defined as the amount of enzyme required to produce 1 µg of acid-soluble total nucleotide in 1 minute at 37°C.

Unit Assay Conditions: 1X Mung Bean Nuclease Reaction Buffer and 0.5 mg/ml denatured calf thymus DNA as an enzyme substrate.

Removal of Single-Stranded Extensions:

1. Suspend DNA (0.1 µg/µl) in 1X Mung Bean Nuclease Reaction Buffer or 1X NEBuffers 1.1, 2.1 or CutSmart.
2. Add 1.0 unit of Mung Bean Nuclease per µg DNA.
3. Incubate at 30°C for 30 minutes.
4. Inactivate the enzyme by phenol/chloroform extraction or by addition of SDS to 0.01%.
5. Recover the DNA by ethanol precipitation.

1X Mung Bean Nuclease Reaction Buffer:

50 mM sodium acetate
30 mM NaCl
1 mM ZnSO₄
pH 5.0 @ 25°C
Also active in NEBuffers 1.1, 2.1 or CutSmart®.

Unit Definition: One unit is defined as the amount of enzyme required to produce 1 µg of acid-soluble total nucleotide in 1 minute at 37°C.

Unit Assay Conditions: 1X Mung Bean Nuclease Reaction Buffer and 0.5 mg/ml denatured calf thymus DNA as an enzyme substrate.

Removal of Single-Stranded Extensions:

1. Suspend DNA (0.1 µg/µl) in 1X Mung Bean Nuclease Reaction Buffer or 1X NEBuffers 1.1, 2.1 or CutSmart.
2. Add 1.0 unit of Mung Bean Nuclease per µg DNA.
3. Incubate at 30°C for 30 minutes.
4. Inactivate the enzyme by phenol/chloroform extraction or by addition of SDS to 0.01%.
5. Recover the DNA by ethanol precipitation.

Quality Assurance: Purified free of double-strand exonuclease contamination.

Quality Control Assays

16 µg of Hae III digested φX174 DNA was incubated with 10 units of Mung Bean Nuclease in a 400 µl volume of 1X NEBuffer 2 for 30 minutes at 30°C. The DNA was then precipitated, ligated with T4 DNA Ligase and recut. 90% of the DNA fragments treated with Mung Bean Nuclease were ligated and of those 95% were recut with Hae III.

References:

1. Kowalski, D. et al. (1976) *Biochemistry* 15, 4457-4463.
2. McCutchan, T.F. et al. (1984) *Science* 225, 626-628.



NEW ENGLAND BIOLABS® and CUTSMART® are registered trademarks of New England Biolabs, Inc.

This product is intended for research purposes only. This product is not intended to be used for therapeutic or diagnostic purposes in humans or animals.

CERTIFICATE OF ANALYSIS

Mung Bean Nuclease



1-800-632-7799
info@neb.com
www.neb.com



M0250S 025160818081

M0250S

1,500 units **Lot: 0251608** **Exp: 8/18**
10,000 U/ml **Store at -20°C**

Description: A single-strand specific DNA and RNA endonuclease which will degrade single-stranded extensions from the ends of DNA and RNA molecules, leaving blunt, ligatable ends.

Source: Mung bean sprouts

Molecular Weight: 39 kDa

Supplied in: 10 mM sodium acetate (pH 5.0)
0.1 mM zinc acetate, 1 mM cysteine, 0.001%
Triton X-100 and 50% glycerol.

Quality Assurance: Purified free of double-strand exonuclease contamination.

Quality Control Assays

16 µg of Hae III digested φX174 DNA was incubated with 10 units of Mung Bean Nuclease in a 400 µl volume of 1X NEBuffer 2 for 30 minutes at 30°C. The DNA was then precipitated, ligated with T4 DNA Ligase and recut. 90% of the DNA fragments treated with Mung Bean Nuclease were ligated and of those 95% were recut with Hae III.

References:

1. Kowalski, D. et al. (1976) *Biochemistry* 15, 4457-4463.
2. McCutchan, T.F. et al. (1984) *Science* 225, 626-628.



NEW ENGLAND BIOLABS® and CUTSMART® are registered trademarks of New England Biolabs, Inc.

This product is intended for research purposes only. This product is not intended to be used for therapeutic or diagnostic purposes in humans or animals.

CERTIFICATE OF ANALYSIS