

Deep Vent_R[™] (exo⁻) DNA Polymerase



M0259S 004121214121



1-800-632-7799
info@neb.com
www.neb.com

M0259S



200 units 2,000 U/ml Lot: 0041212
RECOMBINANT Store at -20°C Exp: 12/14

Description: Deep Vent_R (exo⁻) DNA Polymerase has been genetically engineered to eliminate the 3' 5' proofreading exonuclease activity associated with Deep Vent_R DNA Polymerase. Deep Vent_R (exo⁻) DNA Polymerase is even more stable than Vent_R[™] (exo⁻) DNA Polymerase with a half-life of 23 hours at 95°C and 8 hours at 100°C. Both Deep Vent_R (exo⁻) and Vent_R (exo⁻) DNA Polymerase are suitable for primer extensions and high temperature (72°C) DNA sequencing.

Source: An E. coli strain that carries the Deep Vent (D141A / E143A) DNA Polymerase gene, a genetically engineered form of the native DNA polymerase from Pyrococcus species GB-D. The native organism was isolated from a submarine thermal vent at 2,010 meters (1) and is able to grow at temperatures as high as 104°C.

Applications:

- PCR
- Primer extension

Supplied in: 100 mM KCl, 0.1 mM EDTA, 10 mM Tris-HCl (pH 7.4), 1 mM dithiothreitol, 0.1% Triton[®] X-100 and 50% glycerol.

Reagents Supplied with Enzyme:

10X ThermoPol[®] Reaction Buffer
100 mM MgSO₄.

Reaction Conditions: 1X ThermoPol Reaction Buffer, with or without additional MgSO₄, DNA template, dNTPs, primer and 2-4 units polymerase in a final volume of 100 µl.

1X ThermoPol Reaction Buffer:

20 mM Tris-HCl
10 mM (NH₄)₂SO₄
10 mM KCl
2 mM MgSO₄
0.1% Triton X-100
pH 8.8 @ 25°C

Unit Definition: One unit is defined as the amount of enzyme that will incorporate 10 nmol of dNTP into acid insoluble material in 30 minutes at 75°C.

Unit Assay Conditions: 1X ThermoPol Reaction Buffer, 200 µM each dNTP including [³H]-dTTP, 200 µg/ml activated calf thymus DNA.

Heat Inactivation: No

Quality Control Assays

Exonuclease Activity: Incubation of a 50 µl reaction in ThermoPol Reaction Buffer containing a minimum of 20 units of Deep Vent_R (exo⁻) DNA Polymerase and 1 µg of a mixture of single and double-stranded [³H] E. coli DNA for 4 hours at either 37°C or 75°C releases < 0.1% of the total radioactivity.

Endonuclease Activity: Incubation of a 50 µl reaction in ThermoPol Reaction Buffer containing a minimum of 20 units of Deep Vent_R (exo⁻) DNA Polymerase with 1 µg of supercoiled φX174 DNA for 4 hours at either 37°C or 75°C results in < 10% conversion to the nicked form as determined by agarose gel electrophoresis.

Physical Purity: Purified to > 95% homogeneity as determined by SDS-PAGE analysis using Coomassie Blue detection.

Calculated Half-lives at 95°C:

Deep Vent _R DNA Polymerase	23 hours
Vent _R DNA Polymerase	6.7 hours
Taq DNA Polymerase	1.6 hours

(See other side)

CERTIFICATE OF ANALYSIS



1-800-632-7799
info@neb.com
www.neb.com

References:

1. Jannasch, H. W. et al. (1992) Applied Environ. Microbiol. 58, 3472–3481.

Companion Products Sold Separately:

Magnesium Sulfate (MgSO ₄) Solution #B1003S	6.0 ml
Diluent D #B8004S	4.0 ml
BSA #B9001S	6.0 ml
ThermoPol Reaction Buffer Pack #B9004S	6.0 ml
ThermoPol II (Mg-free) Reaction Buffer Pack #B9005S	6.0 ml
ThermoPol DF (Detergent-free) Reaction Buffer Pack #B9013S	6.0 ml
Deoxynucleotide Solution Set #N0446S	25 µmol each
Deoxynucleotide Solution Mix #N0447S	8 µmol each
#N0447L	40 µmol each

Using NEB Thermophilic DNA Polymerases to Extend a Primer

General Approach—Setting up a Primer Extension Reaction or a PCR Reaction: Basic reaction conditions are 1X ThermoPol Reaction Buffer, DNA template, DNA polymerase, 1–6 mM MgSO₄ (see suggested initial conditions), 200–400 µM each dNTP and 0.4 µM primer.

The three most important variables to optimize are the amount of polymerase, the annealing temperature for the primer and the magnesium level. Each new primer: template may require reoptimization.

Enzyme Amount: It is important to use the optimal amount of enzyme, especially with the proofreading DNA polymerases. Start with 1 unit/100 µl reaction volume for proofreading DNA polymerases or 4 units/100 µl reaction volume for exo⁻ derivatives (for different reaction volumes adjust this ratio accordingly). In general, lower DNA template concentrations in a primer extension reaction necessitate using the lower amount of DNA polymerase within the recommended range.

Recommended ranges are 1–2 units per 100 µl reaction volume for the Vent_r and Deep Vent_r DNA polymerases, and 2–4 units for the Vent_r (exo⁻) and Deep Vent_r (exo⁻) DNA Polymerases.

Annealing Temperature: The optimal annealing temperature for the primer can usually be predicted from any of several standard methods of calculation. If this temperature does not give satisfactory results, the annealing temperature should be examined in 3°C increments. We recommend using NEB's T_m Calculator, available at www.neb.com/TmCalculator to determine appropriate annealing temperatures for PCR.

In general, the Vent_r and Deep Vent_r DNA polymerases use annealing temperatures that tend to be the same, or higher, than annealing temperatures used by other DNA polymerases. (Different annealing temperatures may be required by different polymerases, perhaps due to differences in the K_m for binding DNA).

Magnesium Concentration: The optimal magnesium concentration is usually 2, 4 or 6 mM. If EDTA is present at significant levels in DNA added to your reaction, the test range may

need to be extended higher. For Vent_r and Deep Vent_r DNA Polymerases, primer extensions longer than 2 kb almost always require magnesium levels higher than 2 mM, while for primer extensions shorter than 2 kb, there is no correlation between length and optimum magnesium concentration.

 Annealing temperature



Patents/Disclaimer: Some applications in which this product can be used may be covered by patents issued and applicable in the United States and certain other countries. Because purchase of this product does not include a license to perform any patented application, users of this product may be required to obtain a patent license depending upon the particular application in which the product is used.

NEW ENGLAND BIOLABS', VENT[™] and THERMOPOL[®] are registered trademarks of New England Biolabs, Inc.

DEEP VENT[™] is a trademark of New England Biolabs, Inc.

TRITON[™] is a registered trademark of Union Carbide Corporation.