

Lambda Exonuclease



1-800-632-7799
info@neb.com
www.neb.com



M0262S 008160818081

M0262S



1,000 units **5,000 U/ml** **Lot: 0081608**
RECOMBINANT **Store at -20°C** **Exp: 8/18**

Description: A highly processive enzyme that acts in the 5' to 3' direction, catalyzing the removal of 5' mononucleotides from duplex DNA. The preferred substrate is 5'-phosphorylated double stranded DNA, although it will also degrade single-stranded and non-phosphorylated substrates at a greatly reduced rate. Lambda Exonuclease is unable to initiate DNA digestion at nicks or gaps (1).

Source: A genetic fusion of the *E. coli* Lambda Exonuclease gene with the gene encoding maltose binding protein (MBP). Following affinity chromatography, Lambda Exonuclease is cleaved from the fusion construct and purified away from MBP.

Supplied in: 50 mM NaCl, 25 mM Tris-HCl (pH 8.0), 0.1 mM EDTA, 1 mM dithiothreitol and 50% glycerol.

Reagents Supplied with Enzyme:
10X Lambda Exonuclease Reaction Buffer.

Reaction Conditions:
1X Lambda Exonuclease Reaction Buffer.
Incubate at 37°C.

1X Lambda Exonuclease Reaction Buffer:
67 mM Glycine-KOH
2.5 mM MgCl₂
50 µg/ml BSA
(pH 9.4 @ 25°C)

Unit Definition: One unit is defined as the amount of enzyme required to produce 10 nmol of acid soluble deoxyribonucleotide from double-stranded substrate in a total reaction volume of 50 µl in 30 minutes at 37°C.

Unit Assay Conditions: 67 mM Glycine-KOH (pH 9.4), 2.5 mM MgCl₂, 50 µg/ml BSA and 1 µg sonicated duplex ³H DNA.

Heat Inactivation: 75°C for 10 minutes.

Quality Control Assays

Endonuclease Activity: Incubation of 200 units of Lambda Exonuclease with 1 µg φX174 RF I DNA for 4 hours at 37°C in 50 µl reaction buffer resulted in < 10% conversion to RF II.

Quality Assurance: Purified free of contaminating endonucleases and exonucleases.

Note: 5'-OH ends are digested 20X slower than 5'-PO₄ ends. Single-strand is digested 100X slower than double-strand DNA (1).

Reference:

1. Little, J.W. (1981) *Gene Amplification and Analysis 2*, 135-145.



NEW ENGLAND BIOLABS® is a registered trademark of New England Biolabs, Inc.

This product is intended for research purposes only. This product is not intended to be used for therapeutic or diagnostic purposes in humans or animals.

CERTIFICATE OF ANALYSIS

Lambda Exonuclease



1-800-632-7799
info@neb.com
www.neb.com



M0262S 008160818081

M0262S



1,000 units **5,000 U/ml** **Lot: 0081608**
RECOMBINANT **Store at -20°C** **Exp: 8/18**

Description: A highly processive enzyme that acts in the 5' to 3' direction, catalyzing the removal of 5' mononucleotides from duplex DNA. The preferred substrate is 5'-phosphorylated double stranded DNA, although it will also degrade single-stranded and non-phosphorylated substrates at a greatly reduced rate. Lambda Exonuclease is unable to initiate DNA digestion at nicks or gaps (1).

Source: A genetic fusion of the *E. coli* Lambda Exonuclease gene with the gene encoding maltose binding protein (MBP). Following affinity chromatography, Lambda Exonuclease is cleaved from the fusion construct and purified away from MBP.

Supplied in: 50 mM NaCl, 25 mM Tris-HCl (pH 8.0), 0.1 mM EDTA, 1 mM dithiothreitol and 50% glycerol.

Reagents Supplied with Enzyme:
10X Lambda Exonuclease Reaction Buffer.

Reaction Conditions:
1X Lambda Exonuclease Reaction Buffer.
Incubate at 37°C.

1X Lambda Exonuclease Reaction Buffer:
67 mM Glycine-KOH
2.5 mM MgCl₂
50 µg/ml BSA
(pH 9.4 @ 25°C)

Unit Definition: One unit is defined as the amount of enzyme required to produce 10 nmol of acid soluble deoxyribonucleotide from double-stranded substrate in a total reaction volume of 50 µl in 30 minutes at 37°C.

Unit Assay Conditions: 67 mM Glycine-KOH (pH 9.4), 2.5 mM MgCl₂, 50 µg/ml BSA and 1 µg sonicated duplex ³H DNA.

Heat Inactivation: 75°C for 10 minutes.

Quality Control Assays

Endonuclease Activity: Incubation of 200 units of Lambda Exonuclease with 1 µg φX174 RF I DNA for 4 hours at 37°C in 50 µl reaction buffer resulted in < 10% conversion to RF II.

Quality Assurance: Purified free of contaminating endonucleases and exonucleases.

Note: 5'-OH ends are digested 20X slower than 5'-PO₄ ends. Single-strand is digested 100X slower than double-strand DNA (1).

Reference:

1. Little, J.W. (1981) *Gene Amplification and Analysis 2*, 135-145.



NEW ENGLAND BIOLABS® is a registered trademark of New England Biolabs, Inc.

This product is intended for research purposes only. This product is not intended to be used for therapeutic or diagnostic purposes in humans or animals.

CERTIFICATE OF ANALYSIS