

# Taq 2X Master Mix



1-800-632-7799  
info@neb.com  
www.neb.com



## M0270S



100 reactions Lot: 0191212 Exp: 6/14  
RECOMBINANT Store at -20°C

Description: Taq DNA Polymerase is a thermostable DNA polymerase that possesses a 5' 3' polymerase activity (1,2,3) and a 5' flap endonuclease activity (4,5).

Taq 2X Master Mix is an optimized ready-to-use solution containing Taq DNA Polymerase, dNTPs, MgCl<sub>2</sub>, KCl and stabilizers. It is ideally suited to routine PCR applications from templates including pure DNA solutions, bacterial colonies, and cDNA products. It can amplify up to 4 kb from complex genomic DNA or up to 5 kb from lambda DNA.

Source: An E. coli strain that carries the Taq DNA Polymerase gene from *Thermus aquaticus* YT-1

- Application:
- PCR
  - Primer Extension
  - Colony PCR

Reagents Supplied with Enzyme:  
25 mM MgCl<sub>2</sub>

Reaction Conditions: 1X Taq Master Mix, DNA template and primers in a total reaction volume of 25 or 50 µl.

Taq 1X Master Mix:  
10 mM Tris-HCl (pH 8.6, @ 25°C)  
50 mM KCl  
1.5 mM MgCl<sub>2</sub>  
25 units/ml Taq DNA polymerase  
0.2 mM dNTPs each  
5% glycerol  
0.08% IGEPAL® CA-630  
0.05% Tween® 20

Unit Definition: One unit is defined as the amount of enzyme that will incorporate 10 nmol of dNTP into acid insoluble material in 30 minutes at 75°C.

Unit Assay Conditions: 1X ThermoPol® Reaction Buffer, 200 µM dNTPs including [<sup>3</sup>H]-dTTP and 200 µg/ml activated Calf Thymus DNA.

Heat Inactivation: No

### Quality Control Assays

5 kb Lambda PCR: 25 cycles of PCR amplification of 5 ng Lambda DNA, 0.2 µM primers and 1X Taq Master Mix results in the expected 5 kb product.

3' 5' Exonuclease Activity: Incubation of a 20 µl reaction in ThermoPol Reaction Buffer containing a minimum of 20 units of Taq DNA Polymerase with 10 nM fluorescent internally labeled oligonucleotide for 30 minutes at either 37°C or 75°C yields no detectable 3' 5' degradation as determined by capillary electrophoresis.

Endonuclease Activity: Incubation of a 50 µl reaction in ThermoPol Reaction Buffer containing a minimum of 20 units of Taq DNA Polymerase with 1 µg of supercoiled φX174 DNA for 4 hours at 75°C results in < 10% conversion to the nicked form as determined by agarose gel electrophoresis.

### PCR

The Polymerase Chain Reaction (PCR) is a powerful and sensitive technique for DNA amplification (6). Taq DNA Polymerase is an enzyme widely used in PCR (7). The following guidelines are provided to ensure successful PCR using New England Biolabs' Taq 2X Master Mix. These guidelines cover routine PCR reactions. Amplification of templates with high GC content, high secondary structure, low template concentrations, or amplicons greater than 5 kb may require further optimization.

Reaction setup:

We recommend assembling all reaction components on ice and quickly transferring the reactions to a thermocycler preheated to the denaturation temperature (95°C).

COMPONENT	25 µl REACTION	50 µl REACTION	FINAL CONCENTRATION
10 µM Forward Primer	0.5 µl	1 µl	0.2 µM (0.05–1 µM)
10 µM Reverse Primer	0.5 µl	1 µl	0.2 µM (0.05–1 µM)
Template DNA	variable	variable	<1,000 ng
Taq 2X Master Mix	12.5 µl	25 µl	1X
Nuclease-Free Water	to 25 µl	to 50 µl	

Notes: Gently mix the reaction. Collect all liquid to the bottom of the tube by a quick spin if necessary. Overlay the sample with mineral oil if using a PCR machine without a heated lid.

Transfer PCR tubes from ice to a PCR machine with the block preheated to 95°C and begin thermocycling:

Thermocycling conditions for a routine PCR:

STEP	TEMP	TIME
Initial Denaturation	95°C	30 seconds
30 Cycles	95°C	15–30 seconds
	45–68°C	15–60 seconds
	68°C	1 minute/kb
Final Extension	68°C	5 minutes
Hold	4–10°C	

General Guidelines:

1. Template:  
Use of high quality, purified DNA templates greatly enhances the success of PCR reactions. Recommended amounts of DNA template for a 50 µl reaction are as follows:

DNA	AMOUNT
Genomic	1 ng–1 µg
Plasmid or Viral	1 pg–1 ng

2. Primers:  
Oligonucleotide primers are generally 20–40 nucleotides in length and ideally have a GC content of 40–60%. Computer programs such as Primer3 (<http://frodo.wi.mit.edu/primer3>) can be used to design or analyze primers. The final concentration of each primer in a PCR reaction may be 0.05–1 µM, typically 0.1–0.5 µM.

3. Mg<sup>++</sup> and additives:  
Mg<sup>++</sup> concentration of 1.5–2.0 mM is optimal for most PCR products generated with Taq DNA Polymerase. The final Mg<sup>++</sup> concentration in 1X Taq Master Mix is 1.5 mM. This supports satisfactory amplification of most amplicons. However, Mg<sup>++</sup> can be further optimized in 0.5 or 1.0 mM increments using MgCl<sub>2</sub>.

Amplification of some difficult targets, like GC-rich sequences, may be improved with additives, such as DMSO (8) or formamide (9).

4. Denaturation:  
An initial denaturation of 30 seconds at 95°C is sufficient for most amplicons from pure DNA templates. For difficult templates such as GC-rich sequences, a longer denaturation of 2–4 minutes at 95°C is recommended prior to PCR cycling to fully denature the template. With colony PCR, an initial 5 minute denaturation at 95°C is recommended.

During thermocycling a 15–30 second denaturation at 95°C is recommended.

5. Annealing:  
The annealing step is typically 15–60 seconds. Annealing temperature is based on the T<sub>m</sub> of the primer pair and is typically 45–68°C. Annealing temperatures can be optimized by doing a temperature gradient PCR starting 5°C below the calculated T<sub>m</sub>. We recommend using NEB's T<sub>m</sub> Calculator, available at [www.neb.com/TmCalculator](http://www.neb.com/TmCalculator) to determine appropriate annealing temperatures for PCR.

When primers with annealing temperatures above 60°C are used, a 2-step PCR protocol is possible (see #8).

6. Extension:  
The recommended extension temperature is 68°C. Extension times are generally 1 minute per kb. A final extension of 5 minutes at 68°C is recommended.
7. Cycle number:  
Generally, 25–35 cycles yields sufficient product. Up to 45 cycles may be required to detect low-copy-number targets.

(see other side)

8. 2-step PCR:  
When primers with annealing temperatures above 60°C are used, a 2-step thermocycling protocol is possible.

Thermocycling conditions for a routine 2-step PCR:

STEP	TEMP	TIME
Initial Denaturation	95°C	30 seconds
30 Cycles	95°C	15–30 seconds
	60–68°C	1 minute/kb
Final Extension	60–68°C	5 minutes
Hold	4–10°C	

11. PCR product:  
The PCR products generated using Taq DNA Polymerase contain dA overhangs at the 3'–end; therefore the PCR products can be ligated to dT/dU-overhang vectors.

Notes: Taq 2X Master Mix is stable for fifteen freeze-thaw cycles when stored at –20°C

Taq 2X Master Mix is also stable for three months at 4°C, so for frequent use, an aliquot may be kept at 4°C.

#### References:

- Chien, A., Edgar, D.B. and Trela, J.M. (1976) *J. Bact.*, 127, 1550–1557.
- Kaledin, A.S., Sliusarenko, A.G. and Gorodetskii, S.I. (1980) *Biokhimiya*, 45, 644–651.
- Lawyer, F.C. et al. (1993) *PCR Methods and Appl.*, 2, 275–287.
- Longley, M.J., Bennett, S.E. and Mosbaugh D.W. (1990) *Nucleic Acids Res.*, 18, 7317–7322.
- Lyamichev, V., Brow, M.A. and Dahlberg, J.E. (1993) *Science*, 260, 778–783.
- Saiki R.K. et al. (1985) *Science*, 230, 1350–1354.
- Powell, L.M. et al. (1987) *Cell*, 50, 831–840.
- Sun, Y., Hegamyer, G. and Colburn, N. (1993) *Biotechniques*, 15, 372–374.
- Sarkar, G., Kapelner, S. and Sommer, S.S. (1990) *Nucleic Acids Res.*, 18, 7465.

Companion Products Sold Separately:  
Magnesium Chloride (MgCl<sub>2</sub>) Solution  
#B9021S 6.0 ml

 Annealing temperature



NEW ENGLAND BIOLABS® and THERMOPOL® are registered trademarks of New England Biolabs, Inc.

IGEPAL® is a registered trademark of Rhodia Operations.

TWEEN® is a registered trademark of Uniqema Americas LLC.