

Afu Uracil-DNA Glycosylase (UDG)



1-800-632-7799
info@neb.com
www.neb.com



M0279S 003151217121

M0279S



200 units 2,000 U/ml Lot: 0031512

RECOMBINANT Store at -20°C Exp: 12/17

Description: A thermostable homolog of the *E. coli* Uracil-DNA Glycosylase (UDG) (1,2) from *Archaeoglobus fulgidus*. *Afu* UDG catalyzes the release of free uracil from uracil-containing DNA. *Afu* UDG efficiently hydrolyzes uracil from single-stranded or double-stranded DNA.

Source: Cloned and overexpressed in *E. coli*.

Supplied in: 50 mM KCl, 10 mM Tris-HCl (pH 7.4 @ 25°C), 0.1 mM EDTA, 1 mM DTT, 0.1 mg/ml BSA and 50% glycerol.

New Reaction Buffer

Reagents Supplied with Enzyme:
10X ThermoPol® II (Mg-free) Reaction Buffer.

Reaction Conditions: 1X ThermoPol II (Mg-free) Reaction Buffer. Incubate at 65°C.

1X ThermoPol II (Mg-free) Reaction Buffer:
20 mM Tris-HCl
10 mM KCl
10 mM (NH₄)₂SO₄
0.1% Triton X-100
(pH 8.8 @ 25°C).

Unit Definition: One unit is defined as the amount of enzyme that catalyzes the release of 60 pmol of uracil per minute from double-stranded, uracil-containing DNA. Activity is measured by release of [³H]-uracil in a 50 µl reaction containing 0.2 µg DNA (10⁴-10⁵ cpm/µg) in 30 minutes at 65°C.

Unit Assay Conditions: 1X ThermoPol II (Mg-free) Reaction Buffer, 1 unit of *Afu* Uracil-DNA Glycosylase, 0.2 µg ³H-uracil DNA (10⁴-10⁵ cpm/µg) for 30 minutes at 65°C in a total reaction volume of 50 µl.

Reagents Supplied with Enzyme:
10X ThermoPol® II (Mg-free) Reaction Buffer.

Reaction Conditions: 1X ThermoPol II (Mg-free) Reaction Buffer. Incubate at 65°C.

1X ThermoPol II (Mg-free) Reaction Buffer:
20 mM Tris-HCl
10 mM KCl
10 mM (NH₄)₂SO₄
0.1% Triton X-100
(pH 8.8 @ 25°C).

Unit Definition: One unit is defined as the amount of enzyme that catalyzes the release of 60 pmol of uracil per minute from double-stranded, uracil-containing DNA. Activity is measured by release of [³H]-uracil in a 50 µl reaction containing 0.2 µg DNA (10⁴-10⁵ cpm/µg) in 30 minutes at 65°C.

Unit Assay Conditions: 1X ThermoPol II (Mg-free) Reaction Buffer, 1 unit of *Afu* Uracil-DNA Glycosylase, 0.2 µg ³H-uracil DNA (10⁴-10⁵ cpm/µg) for 30 minutes at 65°C in a total reaction volume of 50 µl.

Molecular Weight: 22,717 daltons.

Specific Activity: 96 units/µg

Quality Control Assays

16-Hour Incubation: A 50 µl reaction containing 1 µg of λ phage DNA in ThermoPol Reaction Buffer and 50 units of *Afu* UDG showed no degradation following overnight incubation at 65°C.

Exonuclease Activity: Incubation of 50 units for 4 hours at 65°C in 50 µl of ThermoPol Reaction Buffer with 1 µg ³H DNA (10⁵ cpm/µg) released < 0.1% radioactivity.

Endonuclease Activity: Incubation of 5 units of *Afu* UDG with 1 µg of φX174 RF I DNA for 4 hours at 65°C in 50 µl ThermoPol Reaction Buffer resulted in < 20% conversion to RF II.

Heat Inactivation: No

Note: *Afu* UDG retains 50% activity in the presence of 150 mM NaCl. *Afu* UDG retains less than 1% activity after boiling for 30 minutes in standard reaction conditions.

Molecular Weight: 22,717 daltons.

Specific Activity: 96 units/µg

Quality Control Assays

16-Hour Incubation: A 50 µl reaction containing 1 µg of λ phage DNA in ThermoPol Reaction Buffer and 50 units of *Afu* UDG showed no degradation following overnight incubation at 65°C.

Exonuclease Activity: Incubation of 50 units for 4 hours at 65°C in 50 µl of ThermoPol Reaction Buffer with 1 µg ³H DNA (10⁵ cpm/µg) released < 0.1% radioactivity.

Endonuclease Activity: Incubation of 5 units of *Afu* UDG with 1 µg of φX174 RF I DNA for 4 hours at 65°C in 50 µl ThermoPol Reaction Buffer resulted in < 20% conversion to RF II.

Heat Inactivation: No

Note: *Afu* UDG retains 50% activity in the presence of 150 mM NaCl. *Afu* UDG retains less than 1% activity after boiling for 30 minutes in standard reaction conditions.

Under standard reaction conditions, uracil glycosylase inhibitor (UGI) does not inhibit *Afu* UDG.

References:

1. Sandigursky, M. and Franklin, W. A. (1999) *Current Biology* 9, 531-534.
2. Sandigursky, M. and Franklin, W. A. (2000) *J. Biol. Chem.* 275, 19146-19149.



NEW ENGLAND BIOLABS® and THERMOPOL® are registered trademarks of New England Biolabs, Inc.

This product is intended for research purposes only. This product is not intended to be used for therapeutic or diagnostic purposes in humans or animals.

CERTIFICATE OF ANALYSIS

Afu Uracil-DNA Glycosylase (UDG)



1-800-632-7799
info@neb.com
www.neb.com



M0279S 003151217121

M0279S



200 units 2,000 U/ml Lot: 0031512

RECOMBINANT Store at -20°C Exp: 12/17

Description: A thermostable homolog of the *E. coli* Uracil-DNA Glycosylase (UDG) (1,2) from *Archaeoglobus fulgidus*. *Afu* UDG catalyzes the release of free uracil from uracil-containing DNA. *Afu* UDG efficiently hydrolyzes uracil from single-stranded or double-stranded DNA.

Source: Cloned and overexpressed in *E. coli*.

Supplied in: 50 mM KCl, 10 mM Tris-HCl (pH 7.4 @ 25°C), 0.1 mM EDTA, 1 mM DTT, 0.1 mg/ml BSA and 50% glycerol.

New Reaction Buffer

Under standard reaction conditions, uracil glycosylase inhibitor (UGI) does not inhibit *Afu* UDG.

References:

1. Sandigursky, M. and Franklin, W. A. (1999) *Current Biology* 9, 531-534.
2. Sandigursky, M. and Franklin, W. A. (2000) *J. Biol. Chem.* 275, 19146-19149.



NEW ENGLAND BIOLABS® and THERMOPOL® are registered trademarks of New England Biolabs, Inc.

This product is intended for research purposes only. This product is not intended to be used for therapeutic or diagnostic purposes in humans or animals.

CERTIFICATE OF ANALYSIS