

# hAAG



1-800-632-7799  
info@neb.com  
www.neb.com



M0313S 002151217121

## M0313S



500 units 10,000 U/ml Lot: 0021512

RECOMBINANT Store at -20°C Exp: 12/17

**Description:** Human Alkyladenine DNA Glycosylase (hAAG) excises alkylated and oxidative DNA damaged sites, including 3-methyladenine, 7-methylguanine, 1,N<sup>6</sup>-ethenoadenine and hypoxanthine. hAAG catalyzes the hydrolysis of the N-glycosidic bond to release the damaged base. hAAG is also known as methylpurine DNA glycosylase (MPG) (1,2,3) or 3-methyladenine-DNA glycosylase (ANPG) (4).

**Source:** An *E. coli* strain which carries the cloned truncated human AAG gene (1)

### Applications:

- Single cell gel electrophoresis (Comet Assay) (5,6,7)
- Alkaline elution (8)
- Alkaline unwinding (9)

Supplied in: 100 mM KCl, 10 mM Tris-HCl (pH7.4), 0.1 mM EDTA, 1 mM dithiothreitol, 0.5% Tween 20, 0.5% NP-40 and 50% glycerol.

### Reagents Supplied with Enzyme:

10X ThermoPol Buffer.

### Reaction Conditions:

1X ThermoPol Buffer. Incubate at 37°C.

### 1X ThermoPol Reaction Buffer:

10 mM KCl  
10 mM (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub>  
20 mM Tris-HCl  
2 mM MgSO<sub>4</sub>  
0.1% Triton X-100  
pH 8.8 @ 25°C

When using a buffer other than the optimal (supplied) NEBuffer, it may be necessary to add more enzyme to achieve complete digestion.

**Unit Definition:** One unit is defined as the amount of enzyme required to create an AP site from 1 pmol of a 34-mer oligonucleotide duplex containing a single deoxyinosine site in a total reaction volume of 10 µl in 1 hour at 37°C.

**Unit Assay Conditions:** 1X ThermoPol Buffer containing 5 pmol of fluorescently labeled oligonucleotide duplex in a total reaction volume of 10 µl.

**Molecular Weight:** 25,752 Daltons

### Quality Control Assays

**Physical Purity:** Purified to > 95% homogeneity as determined by SDS-PAGE analysis using Coomassie Blue detection.

**16-Hour Incubation:** A 50 µl reaction containing 1 µg of λ DNA (HindIII digest) and 100 units of hAAG incubated for 16 hours at 37°C resulted in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.

**Exonuclease Activity:** Incubation of a 50 µl reaction containing 50 units of hAAG in NEBuffer 1 with 1 µg of a mixture of single and double-stranded [<sup>3</sup>H] *E. coli* DNA (10<sup>5</sup> cpm/µg) for 4 hours at 37°C released < 0.1% of the total radioactivity.

**Endonuclease Activity:** Incubation of a 50 µl reaction containing 100 units of hAAG with 1 µg φX174 RF I DNA for 4 hours at 37°C resulted in < 10% conversion to RFLI as determined by agarose gel electrophoresis.

**Heat Inactivation:** 65°C for 20 minutes.

(see other side)

CERTIFICATE OF ANALYSIS

# hAAG



1-800-632-7799  
info@neb.com  
www.neb.com



M0313S 002151217121

## M0313S



500 units 10,000 U/ml Lot: 0021512

RECOMBINANT Store at -20°C Exp: 12/17

**Description:** Human Alkyladenine DNA Glycosylase (hAAG) excises alkylated and oxidative DNA damaged sites, including 3-methyladenine, 7-methylguanine, 1,N<sup>6</sup>-ethenoadenine and hypoxanthine. hAAG catalyzes the hydrolysis of the N-glycosidic bond to release the damaged base. hAAG is also known as methylpurine DNA glycosylase (MPG) (1,2,3) or 3-methyladenine-DNA glycosylase (ANPG) (4).

**Source:** An *E. coli* strain which carries the cloned truncated human AAG gene (1)

### Applications:

- Single cell gel electrophoresis (Comet Assay) (5,6,7)
- Alkaline elution (8)
- Alkaline unwinding (9)

Supplied in: 100 mM KCl, 10 mM Tris-HCl (pH7.4), 0.1 mM EDTA, 1 mM dithiothreitol, 0.5% Tween 20, 0.5% NP-40 and 50% glycerol.

### Reagents Supplied with Enzyme:

10X ThermoPol Buffer.

### Reaction Conditions:

1X ThermoPol Buffer. Incubate at 37°C.

### 1X ThermoPol Reaction Buffer:

10 mM KCl  
10 mM (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub>  
20 mM Tris-HCl  
2 mM MgSO<sub>4</sub>  
0.1% Triton X-100  
pH 8.8 @ 25°C

When using a buffer other than the optimal (supplied) NEBuffer, it may be necessary to add more enzyme to achieve complete digestion.

**Unit Definition:** One unit is defined as the amount of enzyme required to create an AP site from 1 pmol of a 34-mer oligonucleotide duplex containing a single deoxyinosine site in a total reaction volume of 10 µl in 1 hour at 37°C.

**Unit Assay Conditions:** 1X ThermoPol Buffer containing 5 pmol of fluorescently labeled oligonucleotide duplex in a total reaction volume of 10 µl.

**Molecular Weight:** 25,752 Daltons

### Quality Control Assays

**Physical Purity:** Purified to > 95% homogeneity as determined by SDS-PAGE analysis using Coomassie Blue detection.

**16-Hour Incubation:** A 50 µl reaction containing 1 µg of λ DNA (HindIII digest) and 100 units of hAAG incubated for 16 hours at 37°C resulted in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.

**Exonuclease Activity:** Incubation of a 50 µl reaction containing 50 units of hAAG in NEBuffer 1 with 1 µg of a mixture of single and double-stranded [<sup>3</sup>H] *E. coli* DNA (10<sup>5</sup> cpm/µg) for 4 hours at 37°C released < 0.1% of the total radioactivity.

**Endonuclease Activity:** Incubation of a 50 µl reaction containing 100 units of hAAG with 1 µg φX174 RF I DNA for 4 hours at 37°C resulted in < 10% conversion to RFLI as determined by agarose gel electrophoresis.

**Heat Inactivation:** 65°C for 20 minutes.

(see other side)

CERTIFICATE OF ANALYSIS

## References:

1. O'Brien, P. and Ellenberger, T. (2003) *Biochemistry* 42, 12418–12429.
2. Abner, C.W. et al. (2001) *J. Biol. Chem.* 276, 13379–13387.
3. Samson, L. (1991) *Proc. Natl. Acad. Sci.* 88, 9127–9131.
4. Lau, A.Y. et al. (1998) *Cell*, 95, 249–258.
5. Singh, N. et al. (1961) *Experimental Cell Reseach* 175, 184–191.
6. Collins, A. et al. (1993) *Carcinogenesis* 14, 1733–1735.
7. Collins, A. et al. (1996) *Environmental Health Perspectives* 104, 465–469.
8. Pflaum, M. et al. (1998) *Free Rad. Res.* 29, 585–594.
9. Hartwig, A. et al. (1996) *Toxicology Letters* 88, 85–90.



NEW ENGLAND BIOLABS® is a registered trademark of New England Biolabs, Inc.

This product is intended for research purposes only. This product is not intended to be used for therapeutic or diagnostic purposes in humans or animals.

## References:

1. O'Brien, P. and Ellenberger, T. (2003) *Biochemistry* 42, 12418–12429.
2. Abner, C.W. et al. (2001) *J. Biol. Chem.* 276, 13379–13387.
3. Samson, L. (1991) *Proc. Natl. Acad. Sci.* 88, 9127–9131.
4. Lau, A.Y. et al. (1998) *Cell*, 95, 249–258.
5. Singh, N. et al. (1961) *Experimental Cell Reseach* 175, 184–191.
6. Collins, A. et al. (1993) *Carcinogenesis* 14, 1733–1735.
7. Collins, A. et al. (1996) *Environmental Health Perspectives* 104, 465–469.
8. Pflaum, M. et al. (1998) *Free Rad. Res.* 29, 585–594.
9. Hartwig, A. et al. (1996) *Toxicology Letters* 88, 85–90.



NEW ENGLAND BIOLABS® is a registered trademark of New England Biolabs, Inc.

This product is intended for research purposes only. This product is not intended to be used for therapeutic or diagnostic purposes in humans or animals.